



POWER 007000 -
10/50234
INVESTOR IN PEOPLE

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ


REC'D 31 MAR 2003

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




R. Mahoney

Signed

Dated 26 February 2003

An Executive Agency of the Department of Trade and Industry

BEST AVAILABLE COPY



The
Patent
Office

1/77

30JAN02 E691737-1 C69803
P01/7700 0.00-0202025.3

Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference

JBV/PMS/P32975

2. Patent application number

(The Patent Office will fill in this part)

29 JAN 2002

0202025.3

3. Full name, address and postcode of the or of each applicant (underline all surnames)

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

Glaxo Group Ltd
Glaxo Wellcome House, Berkeley Avenue,
Greenford, Middlesex UB6 0NN, Great Britain

United Kingdom

82022 9300

4. Title of the invention

Compounds

5. Name of your agent (if you have one)

"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)

Patents ADP number (if you know it)

Corporate Intellectual Property

GlaxoSmithKline
Two New Horizons Court
BRENTFORD
Middlesex TW8 9EP

7960982002

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or each of these earlier applications and (if you know it) the or each application number

Country	Priority application number (if you know it)	Date of filing (day / month / year)
---------	---	--

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application	Date of filing (day / month / year)
-------------------------------	--

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer yes if:

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is named as an applicant, or
 - c) any named applicant is a corporate body
- See note (d)

Patents Form 1/77

Enter the number of sheets for any of the following items you are filing with this form.
Do not count copies of the same document

Continuation sheets of this form	0
Description	35
Claim(s)	2
Abstract	1
Drawings	0

10. If you are also filing any of the following, state how many against each item.

Priority Documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents
(please specify)

11. We request the grant of a patent on the basis of this application
Signature J B Valentine Date 29-Jan-02
J B Valentine

12. Name and daytime telephone number of person to contact in the United Kingdom
J B Valentine +44 (0)1279 644401

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission unless an application has been filed at least six weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505
- b) Write your answers in capital letters using black ink or you may type them.
- c) If there is not enough space for all relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- d) If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- f) For details of the fee and ways to pay please contact the Patent Office.

Compounds


This invention relates to novel compounds, compositions containing them and their use as antibacterials.

WO99/37635, WO00/21948, WO00/21952, WO00/25227, WO00/43383, WO00/78748, WO01/07432 and WO01/07433 disclose piperidine and piperazine derivatives having antibacterial activity.

WO9717957 discloses piperidyl compounds which are haemoregulatory and stimulate haematopoiesis. JP07179407 discloses piperidyl compounds which are useful for preventing thrombotic diseases, inhibiting tumour metastasis and accelerating wound healing.

We have now found a novel group of aminopiperidines which have antibacterial activity.

This invention provides a compound of formula (I) or a pharmaceutically acceptable derivative thereof:



(I)

wherein:

one of Z^1 , Z^2 , Z^3 , Z^4 and Z^5 is N, one is CR^{1a} and the remainder are CH, or one or two of Z^1 , Z^2 , Z^3 , Z^4 and Z^5 are independently CR^{1a} and the remainder are CH;

R^1 and R^{1a} are independently hydrogen; hydroxy; (C_{1-6}) alkoxy optionally substituted by (C_{1-6}) alkoxy, amino, piperidyl, guanidino or amidino any of which is optionally N-substituted by one or two (C_{1-6}) alkyl, acyl or (C_{1-6}) alkylsulphonyl groups, $CONH_2$, hydroxy, (C_{1-6}) alkylthio, heterocyclylthio, heterocyclyloxy, arylthio, aryloxy, acylthio, acyloxy or (C_{1-6}) alkylsulphonyloxy; (C_{1-6}) alkoxy-substituted (C_{1-6}) alkyl; halogen; (C_{1-6}) alkyl; (C_{1-6}) alkylthio; trifluoromethyl; trifluoromethoxy; nitro; azido; acyl; acyloxy; acylthio; (C_{1-6}) alkylsulphonyl; (C_{1-6}) alkylsulphoxide; arylsulphonyl; arylsulphoxide or an amino, piperidyl, guanidino or amidino group optionally N-substituted by one or two (C_{1-6}) alkyl, acyl or (C_{1-6}) alkylsulphonyl groups;

provided that when Z^1 , Z^2 , Z^3 , Z^4 and Z^5 are CR^{1a} or CH , then R^1 is not hydrogen;

R^2 is hydrogen, or (C_{1-4}) alkyl or (C_{2-4}) alkenyl optionally substituted with 1 to 3 groups selected from:


- 5 amino optionally substituted by one or two (C_{1-4}) alkyl groups; carboxy; (C_{1-4}) alkoxycarbonyl; (C_{1-4}) alkylcarbonyl; (C_{2-4}) alkenyloxycarbonyl; (C_{2-4}) alkenylcarbonyl; aminocarbonyl wherein the amino group is optionally substituted by hydroxy, (C_{1-4}) alkyl, hydroxy (C_{1-4}) alkyl, aminocarbonyl (C_{1-4}) alkyl, (C_{2-4}) alkenyl, (C_{1-4}) alkylsulphonyl, trifluoromethylsulphonyl, (C_{2-4}) alkenylsulphonyl, (C_{1-4}) alkoxycarbonyl, (C_{1-4}) alkylcarbonyl, (C_{2-4}) alkenyloxycarbonyl or (C_{2-4}) alkenylcarbonyl; cyano; tetrazolyl; 2-oxo-oxazolidinyl optionally substituted by R^{10} ; 3-hydroxy-3-cyclobutene-1,2-dione-4-yl; 2,4-thiazolidinedione-5-yl; tetrazol-5-ylaminocarbonyl; 1,2,4-triazol-5-yl optionally substituted by R^{10} ; 5-oxo-1,2,4-oxadiazol-3-yl; halogen; (C_{1-4}) alkylthio; trifluoromethyl; hydroxy optionally substituted by (C_{1-4}) alkyl, (C_{2-4}) alkenyl, (C_{1-4}) alkoxycarbonyl, (C_{1-4}) alkylcarbonyl, (C_{2-4}) alkenyloxycarbonyl, (C_{2-4}) alkenylcarbonyl; oxo; (C_{1-4}) alkylsulphonyl; (C_{2-4}) alkenylsulphonyl; or (C_{1-4}) aminosulphonyl wherein the amino group is optionally substituted by (C_{1-4}) alkyl or (C_{2-4}) alkenyl;
- 20 R^3 is in the 2-, 3- or 4-position and is trifluoromethyl or is in the 2-position and is oxo; or R^3 is in the 3-position and is fluorine or amino wherein the amino group is optionally substituted by: hydroxy; (C_{1-6}) alkylsulphonyl; trifluoromethylsulphonyl; (C_{2-6}) alkenylsulphonyl; (C_{1-6}) alkylcarbonyl; (C_{2-6}) alkenylcarbonyl; (C_{1-6}) alkoxycarbonyl; (C_{2-6}) alkenyloxycarbonyl; (C_{1-6}) alkyl; or (C_{2-6}) alkenyl; wherein a (C_{1-6}) alkyl or (C_{2-6}) alkenyl moiety may be optionally substituted with up to 2 groups R^{12} independently selected from:
- 30 halogen; (C_{1-6}) alkylthio; trifluoromethyl; cyano; carboxy; tetrazolyl; 2-oxo-oxazolidinyl; 3-hydroxy-3-cyclobutene-1,2-dione-4-yl; 2,4-thiazolidinedione-5-yl; tetrazol-5-ylaminocarbonyl; 1,2,4-triazol-5-yl optionally substituted by R^{10} ; or 5-oxo-1,2,4-oxadiazol-3-yl; (C_{1-6}) alkoxycarbonyl; (C_{1-6}) alkylcarbonyl; (C_{2-6}) alkenyloxycarbonyl; (C_{2-6}) alkenylcarbonyl; hydroxy optionally substituted by (C_{1-6}) alkyl, (C_{2-6}) alkenyl, (C_{1-6}) alkylcarbonyl, (C_{2-6}) alkenylcarbonyl or aminocarbonyl wherein the amino group is optionally substituted by (C_{1-6}) alkyl, (C_{2-6}) alkenyl; amino optionally mono- or disubstituted by (C_{1-6}) alkoxycarbonyl, (C_{1-6}) alkylcarbonyl, (C_{2-6}) alkenyloxycarbonyl, (C_{2-6}) alkenylcarbonyl, (C_{1-6}) alkyl, (C_{2-6}) alkenyl, (C_{1-6}) alkylsulphonyl, (C_{2-6}) alkenylsulphonyl or aminocarbonyl wherein the amino group is optionally substituted by (C_{1-6}) alkyl or (C_{2-6}) alkenyl;

in addition when R^3 is disubstituted with a hydroxy or amino containing substituent and carboxy containing substituent these may together form a cyclic ester or amide linkage, respectively;

5 R^4 is a group $-U-R^5$ where

U is selected from CO, SO₂ and CH₂ and

R^5 is an optionally substituted bicyclic carbocyclic or heterocyclic ring system (A):



containing up to four heteroatoms in each ring in which

10 ring (a) is aromatic or non aromatic;

X^1 is C when part of an aromatic ring or CR¹⁴ when part of a non aromatic ring;

X^2 is N, NR¹³, O, S(O)_x, CO or CR¹⁴ when part of an aromatic or non-aromatic ring or may in addition be CR¹⁴R¹⁵ when part of a non aromatic ring;

X^4 is N, NR¹³, O, S(O)_x, CO or CR¹⁴;

15 X^3 and X^5 are independently N or C;

Y^1 is a 1 to 3 atom linker group each atom of which is independently selected from N, NR¹³, O, S(O)_x, CO and CR¹⁴ when part of an aromatic or non-aromatic ring or may additionally be CR¹⁴R¹⁵ when part of a non aromatic ring,

Y^2 is a 2 or 3 atom linker group completing an aromatic ring, each atom of Y^2 being independently selected from N, NR¹³, O, S(O)_x, CO and CR¹⁴;

20 each of R¹⁴ and R¹⁵ is independently selected from: H; (C₁₋₄)alkylthio; halo; carboxy(C₁₋₄)alkyl; halo(C₁₋₄)alkoxy; halo(C₁₋₄)alkyl; (C₁₋₄)alkyl; (C₂₋₄)alkenyl; (C₁₋₄)alkoxycarbonyl; formyl; (C₁₋₄)alkylcarbonyl; (C₂₋₄)alkenyloxycarbonyl; (C₂₋₄)alkenylcarbonyl; (C₁₋₄)alkylcarbonyloxy; (C₁₋₄)alkoxycarbonyl(C₁₋₄)alkyl; hydroxy; 25 hydroxy(C₁₋₄)alkyl; mercapto(C₁₋₄)alkyl; (C₁₋₄)alkoxy; nitro; cyano; carboxy; amino or aminocarbonyl optionally substituted as for corresponding substituents in R³; (C₁₋₄)alkylsulphonyl; (C₂₋₄)alkenylsulphonyl; or aminosulphonyl wherein the amino group is optionally mono- or di-substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl; aryl; aryl(C₁₋₄)alkyl; aryl(C₁₋₄)alkoxy or

30 R¹⁴ and R¹⁵ may together represent oxo;

each R¹³ is independently H; trifluoromethyl; (C₁₋₄)alkyl optionally substituted by hydroxy, (C₁₋₆)alkoxy, (C₁₋₆)alkylthio, carboxy, halo or trifluoromethyl; (C₂₋₄)alkenyl; aryl; aryl (C₁₋₄)alkyl; arylcarbonyl; heteroarylcarbonyl; (C₁₋₄)alkoxycarbonyl; (C₁₋₄)alkylcarbonyl; formyl; (C₁₋₆)alkylsulphonyl; or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₄)alkoxycarbonyl, (C₁₋₄)alkylcarbonyl, (C₂₋

35

4)alkenyloxycarbonyl, (C₂₋₄)alkenylcarbonyl, (C₁₋₄)alkyl or (C₂₋₄)alkenyl and optionally further substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl;

n is 0 or 1;

5

each x is independently 0, 1 or 2

A is NR¹¹, O or CR⁶R⁷ and B is NR¹¹, O, SO₂ or CR⁸R⁹ and wherein:

each of R⁶, R⁷, R⁸ and R⁹ is independently selected from: hydrogen; (C₁₋₆)alkoxy; (C₁₋₆)alkylthio; halo; trifluoromethyl; azido; (C₁₋₆)alkyl; (C₂₋₆)alkenyl; (C₁₋

10

6)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; (C₂₋₆)alkenylcarbonyl; hydroxy, amino or aminocarbonyl optionally substituted as for corresponding substituents in R³; (C₁₋₆)alkylsulphonyl; (C₂₋₆)alkenylsulphonyl; or aminosulphonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl or (C₂₋

15

6)alkenyl; or when n=1 R⁶ and R⁸ together represent a bond and R⁷ and R⁹ are as above defined; or R⁶ and R⁷ or R⁸ and R⁹ together represent oxo;

provided that:

20

when A is NR¹¹, B is not NR¹¹ or O;

when A is CO, B is not CO, O or SO₂;

when n is 0 and A is NR¹¹, CR⁸R⁹ can only be CO;

when A is CR⁶R⁷ and B is SO₂, n is 0;

when n is 0, B is not NR¹¹ or O or R⁸ and R⁹ are not optionally substituted hydroxy or

25

amino;

when A is O, B is not NR¹¹, O, SO₂ or CO and n=1; and

when A-B is CR⁷=CR⁹, n is 1

R¹⁰ is selected from (C₁₋₄)alkyl; (C₂₋₄)alkenyl and aryl any of which may be optionally substituted by a group R¹² as defined above; carboxy; aminocarbonyl wherein the amino group is optionally substituted by hydroxy, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylsulphonyl, trifluoromethylsulphonyl, (C₂₋₆)alkenylsulphonyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl or (C₂₋

30

6)alkenylcarbonyl and optionally further substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl;

35

(C₁₋₆)alkylsulphonyl; trifluoromethylsulphonyl; (C₂₋₆)alkenylsulphonyl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; and (C₂₋

6)alkenylcarbonyl; and

R¹¹ is hydrogen; trifluoromethyl, (C₁₋₆)alkyl; (C₂₋₆)alkenyl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl, (C₂₋₆)alkenylcarbonyl, (C₁₋₆)alkyl or (C₂₋₆)alkenyl and optionally further substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl;

or where one of R³ and R⁶, R⁷, R⁸ or R⁹ contains a carboxy group and the other contains a hydroxy or amino group they may together form a cyclic ester or amide linkage.

This invention also provides a method of treatment of bacterial infections in mammals, particularly in man, which method comprises the administration to a mammal in need of such treatment an effective amount of a compound of formula (I), or a pharmaceutically acceptable derivative thereof.

The invention also provides the use of a compound of formula (I), or a pharmaceutically acceptable derivative thereof, in the manufacture of a medicament for use in the treatment of bacterial infections in mammals.

The invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable derivative thereof, and a pharmaceutically acceptable carrier.

Preferably one of Z¹, Z², Z³, Z⁴ and Z⁵ is N, one is CR^{1a} and the remainder are CH, or one of Z¹, Z², Z³, Z⁴ and Z⁵ is CR^{1a} and the remainder are CH.

Preferably Z⁵ is CH or N, Z³ is CH or CF and Z¹, Z² and Z⁴ are each CH, or Z¹ is N, Z³ is CH or CF and Z², Z⁴ and Z⁵ are each CH.

When R¹ or R^{1a} is substituted alkoxy it is preferably (C₂₋₆)alkoxy substituted by optionally N-substituted amino, guanidino or amidino, or (C₁₋₆)alkoxy substituted by piperidyl. Suitable examples of R¹ and R^{1a} alkoxy include methoxy, trifluoromethoxy, n-propyloxy, iso-butyloxy, aminoethyloxy, aminopropyloxy, aminobutyloxy, aminopentyloxy, guanidinopropyloxy, piperidin-4-ylmethyloxy, phthalimido pentyloxy or 2-aminocarbonylprop-2-oxy.

Preferably R¹ and R^{1a} are independently methoxy, amino(C₃₋₅)alkyloxy, guanidino(C₃₋₅)alkyloxy, piperidyl(C₃₋₅)alkyloxy, nitro or fluoro; R¹ is more preferably methoxy, amino(C₃₋₅)alkyloxy or guanidino(C₃₋₅)alkyloxy. R^{1a} is more preferably H or F. Most preferably R¹ is methoxy. and R^{1a} is H or when Z³ is CR^{1a} it may be C-F.

When Z⁵ is CR^{1a}, R^{1a} is preferably hydrogen, cyano, hydroxymethyl or carboxy, most preferably hydrogen.

Preferably n is 0.

R^2 is preferably hydrogen; (C_{1-4}) alkyl substituted with carboxy, optionally substituted hydroxy, optionally substituted aminocarbonyl, optionally substituted amino or (C_{1-4}) alkoxycarbonyl; or (C_{2-4}) alkenyl substituted with (C_{1-4}) alkoxycarbonyl or carboxy. More preferred groups for R^2 are hydrogen, carboxymethyl, hydroxyethyl, aminocarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylallyl and carboxyallyl, most preferably hydrogen.

Examples of R^3 include CF_3 , fluoro and oxo.

When R^3 is amino it is preferably unsubstituted or substituted by (C_{1-6}) alkyl or (C_{2-6}) alkenyl.

R^3 is preferably in the 3- or 4-position.

When R^3 and R^6 , R^7 , R^8 or R^9 together form a cyclic ester or amide linkage, it is preferred that the resulting ring is 5-7 membered. It is further preferred that the group A or B which does not form the ester or amide linkage is CH_2 .

When A is $CH(OH)$ the *R*-stereochemistry is preferred.

Preferably A is NH , NCH_3 , CH_2 , $CHOH$, $CH(NH_2)$, $C(Me)(OH)$ or $CH(Me)$.

Preferably B is CH_2 or CO .

Preferably A-B is $CHOH-CH_2$, $NR^{11}-CH_2$ or $NR^{11}-CO$.

Particularly preferred are those compounds where $n=0$, A is NH and B is CO , or A is $CHOH$ and B is CH_2 , when more preferably A is the *R*-isomer of $CHOH$.

Preferably R^{11} is hydrogen or (C_{1-4}) alkyl e.g. methyl, more preferably hydrogen.

U is most preferably CH_2 .

Preferably R^5 is an aromatic heterocyclic ring (A) having 1-4 heteroatoms of which one is N or NR^{13} .

Examples of rings (A) include optionally substituted:

(a) and (b) aromatic

1H-pyrrolo[2,3-b]-pyridin-2-yl, 1H-pyrrolo[3,2-b]-pyridin-2-yl, 3H-imidazo[4,5-b]-pyridin-2-yl, 3H-quinazolin-4-one-2-yl, benzimidazol-2-yl, benzo[1,2,3]-thiadiazol-5-yl, benzo[1,2,5]-oxadiazol-5-yl, benzofur-2-yl, benzothiazol-2-yl, benzo[b]thiophen-2-yl, benzoxazol-2-yl, chromen-4-one-3-yl, imidazo[1,2-a]pyridin-2-yl, imidazo-[1,2-a]-pyrimidin-2-yl, indol-2-yl, indol-6-yl, isoquinolin-3-yl, [1,8]-naphthyridine-3-yl, oxazolo[4,5-b]-pyridin-2-yl, quinolin-2-yl, quinolin-3-yl, quinoxalin-2-yl, indan-2-yl, naphthalen-2-yl, 1,3-dioxo-isoindol-2-yl, benzimidazol-2-yl, benzothiophen-2-yl, 1H-benzotriazol-5-yl, 1H-indol-5-yl, 3H-benzooxazol-2-one-6-yl, 3H-benzooxazol-2-thione-6-yl, 3H-benzothiazol-2-one-5-yl, 3H-quinazolin-4-one-2-yl, 3H-quinazolin-4-one-6-yl, 4-oxo-4H-pyrido[1,2-a]pyrimidin-3-yl, benzo[1,2,3]thiadiazol-6-yl, benzo[1,2,5]thiadiazol-5-yl, benzo[1,4]oxazin-2-one-3-yl, benzothiazol-5-yl,

benzothiazol-6-yl, cinnolin-3-yl, imidazo[1,2-a]pyridazin-2-yl, imidazo[1,2-b]pyridazin-2-yl, pyrazolo[1,5-a]pyrazin-2-yl, pyrazolo[1,5-a]pyridin-2-yl, pyrazolo[1,5-a]pyrimidin-6-yl, pyrazolo[5,1-c][1,2,4]triazin-3-yl, pyrido[1,2-a]pyrimidin-4-one-2-yl, pyrido[1,2-a]pyrimidin-4-one-3-yl, quinazolin-2-yl, quinoxalin-6-yl, thiazolo[3,2-a]pyrimidin-5-one-7-yl, thiazolo[5,4-b]pyridin-2-yl, thiazolo[5,4-b]pyridin-6-yl, thieno[3,2-b]pyridin-6-yl, 2H-isoquinolin-1-one-3-yl.

(a) is non aromatic

(2S)-2,3-dihydro-1H-indol-2-yl, (2S)-2,3-dihydro-benzo[1,4]dioxine-2-yl, 3-(R,S)-3,4-dihydro-2H-benzo[1,4]thiazin-3-yl, 3-(R)-(2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl, 2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl, and 3-substituted-3H-quinazolin-4-one-2-yl.

In one aspect, when R¹³ is optionally substituted (C₁₋₄)alkyl, the optional substituent is other than carboxy.

R¹³ is preferably H if in ring (a) or in addition (C₁₋₄)alkyl such as methyl or isopropyl when in ring (b). More preferably, in ring (b) R¹³ is hydrogen when NR¹³ is bonded to X³ and (C₁₋₄)alkyl when NR¹³ is bonded to X⁵.

R¹⁴ and R¹⁵ are preferably independently selected from hydrogen, halo, hydroxy, (C₁₋₄)alkyl, (C₁₋₄)alkoxy, trifluoromethoxy, nitro, cyano, aryl(C₁₋₄)alkoxy and (C₁₋₄)alkylsulphonyl. More preferably R¹⁵ is hydrogen.

More preferably each R¹⁴ is selected from hydrogen, chloro, fluoro, hydroxy, methyl, methoxy, trifluoromethoxy, benzyloxy, nitro, cyano and methylsulphonyl. Most preferably R¹⁴ is selected from hydrogen, hydroxy, fluorine or nitro. Preferably 0-3 groups R¹⁴ are substituents other than hydrogen. Preferably when R¹⁴ is not hydrogen, X⁴ is CR¹⁴ and/or CR¹⁴ is a component of Y².

Most preferred groups R⁵ include 4,6-difluoro-indol-2-yl, 1H-pyrrolo[2,3-b]pyridin-2-yl, 1H-pyrrolo[3,2-b]pyridin-2-yl, 8-hydroxy-quinolin-2-yl, quinoxalin-2-yl, benzimidazol-2-yl, benzo[1,2,3]-thiadiazol-5-yl, benzothiophen-2-yl, 4,6-difluoro-1H-benzimidazol-2-yl, benzothiazole-5-yl and 3-(R)-2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl.

When used herein, the term "alkyl" includes groups having straight and branched chains, for instance, methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl, pentyl and hexyl. The term 'alkenyl' should be interpreted accordingly.

Halo or halogen includes fluoro, chloro, bromo and iodo.

Haloalkyl moieties include 1-3 halogen atoms.

Unless otherwise defined, the term "heterocyclic" as used herein includes optionally substituted aromatic and non-aromatic, single and fused, rings suitably

containing up to four hetero-atoms in each ring selected from oxygen, nitrogen and sulphur, which rings may be unsubstituted or C-substituted by, for example, up to three groups selected from (C₁₋₄)alkylthio; halo; carboxy(C₁₋₄)alkyl; halo(C₁₋₄)alkoxy; halo(C₁₋₄)alkyl; (C₁₋₄)alkyl; (C₂₋₄)alkenyl; (C₁₋₄)alkoxycarbonyl; formyl; (C₁₋₄)alkylcarbonyl; (C₂₋₄)alkenyloxycarbonyl; (C₂₋₄)alkenylcarbonyl; (C₁₋₄)alkylcarbonyloxy; (C₁₋₄)alkoxycarbonyl(C₁₋₄)alkyl; hydroxy; hydroxy(C₁₋₄)alkyl; mercapto(C₁₋₄)alkyl; (C₁₋₄)alkoxy; nitro; cyano, carboxy; amino or aminocarbonyl optionally substituted as for corresponding substituents in R³; (C₁₋₄)alkylsulphonyl; (C₂₋₄)alkenylsulphonyl; or aminosulphonyl wherein the amino group is optionally substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl; optionally substituted aryl, aryl(C₁₋₄)alkyl or aryl(C₁₋₄)alkoxy and oxo groups.

Each heterocyclic ring suitably has from 4 to 7, preferably 5 or 6, ring atoms. A fused heterocyclic ring system may include carbocyclic rings and need include only one heterocyclic ring. Compounds within the invention containing a heterocyclyl group may occur in two or more tautomeric forms depending on the nature of the heterocyclyl group; all such tautomeric forms are included within the scope of the invention.

Where an amino group forms part of a single or fused non-aromatic heterocyclic ring as defined above suitable optional substituents in such substituted amino groups include H; trifluoromethyl; (C₁₋₄)alkyl optionally substituted by hydroxy, (C₁₋₆)alkoxy, (C₁₋₆)alkylthio, halo or trifluoromethyl; (C₂₋₄)alkenyl; aryl; aryl (C₁₋₄)alkyl; (C₁₋₄)alkoxycarbonyl; (C₁₋₄)alkylcarbonyl; formyl; (C₁₋₆)alkylsulphonyl; or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₄)alkoxycarbonyl, (C₁₋₄)alkylcarbonyl, (C₂₋₄)alkenyloxycarbonyl, (C₂₋₄)alkenylcarbonyl, (C₁₋₄)alkyl or (C₂₋₄)alkenyl and optionally further substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl.

When used herein the term "aryl", includes optionally substituted phenyl and naphthyl.

Aryl groups may be optionally substituted with up to five, preferably up to three, groups selected from (C₁₋₄)alkylthio; halo; carboxy(C₁₋₄)alkyl; halo(C₁₋₄)alkoxy; halo(C₁₋₄)alkyl; (C₁₋₄)alkyl; (C₂₋₄)alkenyl; (C₁₋₄)alkoxycarbonyl; formyl; (C₁₋₄)alkylcarbonyl; (C₂₋₄)alkenyloxycarbonyl; (C₂₋₄)alkenylcarbonyl; (C₁₋₄)alkylcarbonyloxy; (C₁₋₄)alkoxycarbonyl(C₁₋₄)alkyl; hydroxy; hydroxy(C₁₋₄)alkyl; mercapto(C₁₋₄)alkyl; (C₁₋₄)alkoxy; nitro; cyano; carboxy; amino or aminocarbonyl optionally substituted as for corresponding substituents in R³; (C₁₋₄)alkylsulphonyl; (C₂₋₄)alkenylsulphonyl; or aminosulphonyl wherein the amino group is optionally substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl; phenyl, phenyl(C₁₋₄)alkyl or phenyl(C₁₋₄)alkoxy

The term "acyl" includes formyl and (C₁₋₆)alkylcarbonyl group.


Some of the compounds of this invention may be crystallised or recrystallised from solvents such as aqueous and organic solvents. In such cases solvates may be formed. This invention includes within its scope stoichiometric solvates including hydrates as well as compounds containing variable amounts of water that may be produced by processes such as lyophilisation.

Since the compounds of formula (I) are intended for use in pharmaceutical compositions it will readily be understood that they are each preferably provided in substantially pure form, for example at least 60% pure, more suitably at least 75% pure and preferably at least 85%, especially at least 98% pure (% are on a weight for weight basis). Impure preparations of the compounds may be used for preparing the more pure forms used in the pharmaceutical compositions; these less pure preparations of the compounds should contain at least 1%, more suitably at least 5% and preferably from 10 to 59% of a compound of the formula (I) or pharmaceutically acceptable derivative thereof.

Particular compounds according to the invention include those mentioned in the examples and their pharmaceutically acceptable derivatives.

Pharmaceutically acceptable derivatives of the above-mentioned compounds of formula (I) include the free base form or their acid addition or quaternary ammonium salts, for example their salts with mineral acids e.g. hydrochloric, hydrobromic, sulphuric nitric or phosphoric acids, or organic acids, e.g. acetic, fumaric, succinic, maleic, citric, benzoic, p-toluenesulphonic, methanesulphonic, naphthalenesulphonic acid or tartaric acids. Compounds of formula (I) may also be prepared as the N-oxide. Compounds of formula (I) having a free carboxy group may also be prepared as an *in vivo* hydrolysable ester. The invention extends to all such derivatives.

Examples of suitable pharmaceutically acceptable *in vivo* hydrolysable ester-forming groups include those forming esters which break down readily in the human body to leave the parent acid or its salt. Suitable groups of this type include those of part formulae (i), (ii), (iii), (iv) and (v):




5

wherein R^a is hydrogen, (C₁₋₆) alkyl, (C₃₋₇) cycloalkyl, methyl, or phenyl, R^b is (C₁₋₆) alkyl, (C₁₋₆) alkoxy, phenyl, benzyl, (C₃₋₇) cycloalkyl, (C₃₋₇) cycloalkyloxy, (C₁₋₆) alkyl (C₃₋₇) cycloalkyl, 1-amino (C₁₋₆) alkyl, or 1-(C₁₋₆ alkyl)amino (C₁₋₆) alkyl; or R^a and R^b together form a 1,2-phenylene group optionally substituted by one or two methoxy groups; R^c represents (C₁₋₆) alkylene optionally substituted with a methyl or ethyl group and R^d and R^e independently represent (C₁₋₆) alkyl; R^f represents (C₁₋₆) alkyl; R^g represents hydrogen or phenyl optionally substituted by up to three groups selected from halogen, (C₁₋₆) alkyl, or (C₁₋₆) alkoxy; Q is oxygen or NH; R^h is hydrogen or (C₁₋₆) alkyl; R^i is hydrogen, (C₁₋₆) alkyl optionally substituted by halogen, (C₂₋₆) alkenyl, (C₁₋₆) alkoxycarbonyl, aryl or heteroaryl; or R^h and R^i together form (C₁₋₆) alkylene; R^j represents hydrogen, (C₁₋₆) alkyl or (C₁₋₆) alkoxycarbonyl; and R^k represents (C₁₋₈) alkyl, (C₁₋₈) alkoxy, (C₁₋₆) alkoxy(C₁₋₆)alkoxy or aryl.

Examples of suitable *in vivo* hydrolysable ester groups include, for example, acyloxy(C₁₋₆)alkyl groups such as acetoxymethyl, pivaloyloxymethyl, α -acetoxylethyl, α -pivaloyloxyethyl, 1-(cyclohexylcarbonyloxy)prop-1-yl, and (1-aminoethyl)carbonyloxymethyl; (C₁₋₆)alkoxycarbonyloxy(C₁₋₆)alkyl groups, such as ethoxycarbonyloxymethyl, α -ethoxycarbonyloxyethyl and propoxycarbonyloxyethyl; di(C₁₋₆)alkylamino(C₁₋₆)alkyl especially di(C₁₋₄)alkylamino(C₁₋₄)alkyl groups such as

dimethylaminomethyl, dimethylaminoethyl, diethylaminomethyl or diethylaminoethyl;
 2-((C₁₋₆)alkoxycarbonyl)-2-(C₂₋₆)alkenyl groups such as
 2-(isobutoxycarbonyl)pent-2-enyl and 2-(ethoxycarbonyl)but-2-enyl; lactone groups such
 as phthalidyl and dimethoxyphthalidyl.

- 5 A further suitable pharmaceutically acceptable *in vivo* hydrolysable ester-forming group is that of the formula:



wherein R^k is hydrogen, C₁₋₆ alkyl or phenyl.


- 10 R is preferably hydrogen.

Compounds of formula (I) may also be prepared as the corresponding N-oxides.


- Certain of the compounds of formula (I) may exist in the form of optical isomers, e.g. diastereoisomers and mixtures of isomers in all ratios, e.g. racemic mixtures. The invention includes all such forms, in particular the pure isomeric forms. For example the
 15 invention includes compound in which an A-B group CH(OH)-CH₂ is in either isomeric configuration, the *R*-isomer is preferred. The different isomeric forms may be separated or resolved one from the other by conventional methods, or any given isomer may be obtained by conventional synthetic methods or by stereospecific or asymmetric syntheses.

- In a further aspect of the invention there is provided a process for preparing
 20 compounds of formula (I), and pharmaceutically acceptable derivatives thereof, which process comprises:

reacting a compound of formula (IV) with a compound of formula (V):



(IV)



(V)


wherein n is as defined in formula (I); Z^{1'}, Z^{2'}, Z^{3'}, Z^{4'}, Z^{5'}, R^{1'} and R^{3'} are Z¹, Z², Z³, Z⁴, Z⁵, R¹ and R³ as defined in formula (I) or groups convertible thereto;

Q^1 is $NR^{2'}R^{4'}$ or a group convertible thereto wherein $R^{2'}$ and $R^{4'}$ are R^2 and R^4 as defined in formula (I) or groups convertible thereto and Q^2 is H or $R^{3'}$ or Q^1 and Q^2 together form an optionally protected oxo group;

and X and Y may be the following combinations:

- 5 (i) X is $A'-COW$, Y is H and n is 0;
- (ii) X is $CR^6=CR^8R^9$, Y is H and n is 0;
- (iii) X is oxirane, Y is H and n is 0;
- (iv) X is $N=C=O$ and Y is H and n is 0;
- (v) one of X and Y is CO_2R^Y and the other is $CH_2CO_2R^X$;
- 10 (vi) X is CHR^6R^7 and Y is $C(=O)R^9$;
- (vii) X is $CR^7=PR^Z_3$ and Y is $C(=O)R^9$ and n=1;
- (viii) X is $C(=O)R^7$ and Y is $CR^9=PR^Z_3$ and n=1;
- (ix) Y is COW and X is $NHR^{11'}$, NCO or $NR^{11'}COW$ and n=0 or 1 or when n=1 X is COW and Y is $NHR^{11'}$, NCO or $NR^{11'}COW$;
- 15 (x) X is $C(O)=R^6$ and Y is $NHR^{11'}$ or X is $NHR^{11'}$ and Y is $C(=O)R^8$ and n=1;
- (xi) X is $NHR^{11'}$ and Y is CR^8R^9W and n=1;
- (xii) X is CR^6R^7W and Y is $NHR^{11'}$ or OH and n=1;
- (xiii) X is $CR^6R^7SO_2W$ and Y is H and n=0;
- (xiv) X is W or OH and Y is CH_2OH and n=1;
- 20 (xv) X is $NHR^{11'}$ and Y is SO_2W or X is $NR^{11'}SO_2W$ and Y is H, and n=0;
- (xvi) X is $NR^{11'}COCH_2W$ or $NR^{11'}SO_2CH_2W$ and Y is H and n=0;
- (xvii) X is W and Y is $CONHR^{11'}$;

- 25 in which W is a leaving group, e.g. halo or imidazolyl; R^X and R^Y are (C_{1-6}) alkyl; R^Z is aryl or (C_{1-6}) alkyl; A' and $NR^{11'}$ are A and NR^{11} as defined in formula (I), or groups convertible thereto; and oxirane is:



wherein R^6 , R^8 and R^9 are as defined in formula (I);

- 30 and thereafter optionally or as necessary converting Q^1 and Q^2 to $NR^{2'}R^{4'}$; converting A' , $Z^{1'}$, $Z^{2'}$, $Z^{3'}$, $Z^{4'}$, $Z^{5'}$, $R^{1'}$, $R^{2'}$, $R^{3'}$, $R^{4'}$ and $NR^{11'}$ to A, Z^1 , Z^2 , Z^3 , Z^4 , Z^5 , R^1 , R^2 , R^3 , R^4 and NR^{11} ; converting A-B to other A-B, interconverting R^1 , R^2 , R^3 and/or R^4 , and/or forming a pharmaceutically acceptable derivative thereof.

- 35 Process variant (i) initially produces compounds of formula (I) wherein A-B is $A'-CO$.

Process variant (ii) initially produces compounds of formula (I) wherein A-B is $\text{CHR}^6\text{-CR}^8\text{R}^9$.

Process variant (iii) initially produces compounds of formula (I) wherein A-B is $\text{CR}^6(\text{OH})\text{-CR}^8\text{R}^9$.

5 Process variant (iv) initially produces compounds of formula (I) where A-B is NH-CO .

Process variant (v) initially produces compounds of formula (I) wherein A-B is CO-CH_2 or $\text{CH}_2\text{-CO}$.

10 Process variant (vi) initially produces compounds of formula (I) wherein A-B is $\text{CR}^6\text{R}^7\text{-CR}^9\text{OH}$.

Process variant (vii) and (viii) initially produce compounds of formula (I) wherein A-B is $\text{CR}^7=\text{CR}^9$.

Process variant (ix) initially produces compounds of formula (I) where A-B is CO-NR^{11} or $\text{NR}^{11}\text{-CO}$.

15 Process variant (x) initially produces compounds of formula (I) wherein A-B is $\text{CHR}^6\text{-NR}^{11}$ or $\text{NR}^{11}\text{-CHR}^8$.

Process variant (xi) initially produces compounds of formula (I) wherein A-B is $\text{NR}^{11'}\text{-CR}^8\text{R}^9$.

20 Process variant (xii) initially produces compounds of formula (I) wherein A-B is $\text{CR}^6\text{R}^7\text{-NR}^{11'}$ or $\text{CR}^6\text{R}^7\text{-O}$.

Process variant (xiii) initially produces compounds of formula (I) where A-B is $\text{CR}^6\text{R}^7\text{-SO}_2$.

Process variant (xiv) initially produces compounds of formula (I) wherein A-B is O-CH_2 .

25 Process variant (xv) initially produces compounds where AB is $\text{NR}^{11}\text{SO}_2$.

Process variant (xvi) initially produces compounds of formula (I) where A-B is $\text{NR}^{11'}\text{-CO}$ or $\text{NR}^{11'}\text{-SO}_2$ and $n=1$.

Process variant (xvii) initially produces compounds of formula (I) where A-B is $\text{NR}^{11'}\text{-CO}$.

30 In process variants (i) and (ix) the reaction is a standard amide or urea formation reaction involving e.g.:

1. Activation of a carboxylic acid (e.g. to an acid chloride, mixed anhydride, active ester, O-acyl-isourea or other species), and treatment with an amine (Ogliaruso, M.A.; Wolfe,

35 J.F. in *The Chemistry of Functional Groups (Ed. Patai, S.) Suppl. B: The Chemistry of Acid Derivatives, Pt. 1* (John Wiley and Sons, 1979), pp 442-8; Beckwith, A.L.J. in *The Chemistry of Functional Groups (Ed. Patai, S.) Suppl. B: The Chemistry of Amides (Ed. Zabricky, J.)* (John Wiley and Sons, 1970), p 73 ff. The acid and amide are preferably

reacted in the presence of an activating agent such as 1-(dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) or 1-hydroxybenzotriazole (HOBT) or O-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU); or

2. The specific methods of:

- 5 a. *in situ* conversion of an acid into the amine component by a modified Curtius reaction procedure (Shioiri, T., Murata, M., Hamada, Y., *Chem. Pharm. Bull.* 1987, 35, 2698)
- b. *in situ* conversion of the acid component into the acid chloride under neutral conditions (Villeneuve, G. B.; Chan, T. H., *Tetrahedron. Lett.* 1997, 38, 6489).

A' may be, for example, protected hydroxymethylene.

- 10 The process variant (ii) is a standard addition reaction using methods well known to those skilled in the art. The process is preferably carried out in a polar organic solvent e.g. acetonitrile in the presence of an organic base e.g. triethylamine.

- In process variant (iii) the coupling may be effected in acetonitrile at room temperature in the presence of one equivalent of lithium perchlorate as catalyst (general
- 15 method of J.E. Chateaneuf *et al.*, *J. Org. Chem.*, 56, 5939-5942, 1991) or more preferably with ytterbium triflate in dichloromethane. In some cases an elevated temperature such as 40 – 70 °C may be beneficial. Alternatively, the piperidine may be treated with a base, such as one equivalent of butyl lithium, and the resulting salt reacted with the oxirane in an inert solvent such as tetrahydrofuran, preferably at an elevated
 - 20 temperature such as 80°C. Use of a chiral epoxide will afford single diastereomers. Alternatively, mixtures of diastereomers may be separated by preparative HPLC or by conventional resolution through crystallisation of salts formed from chiral acids.

- The process variant (iv) is a standard urea formation reaction from the reaction of an isocyanate with an amine and is conducted by methods well known to those skilled in
- 25 the art (for example see March, J; *Advanced Organic Chemistry, Edition 3* (John Wiley and Sons, 1985), p802-3). The process is preferably carried out in a polar solvent such as *N,N*-dimethylformamide.

- In process variant (v) the process is two step: firstly a condensation using a base, preferably sodium hydride or alkoxide, sodamide, alkyl lithium or lithium dialkylamide,
- 30 preferably in an aprotic solvent e.g. ether, THF or benzene; secondly, hydrolysis using an inorganic acid, preferably HCl in aqueous organic solvent at 0-100°C. Analogous routes are described in DE330945, EP31753, EP53964 and H. Sargent, *J. Am. Chem. Soc.* **68**, 2688-2692 (1946). Similar Claisen methodology is described in Soszko *et. al.*, *Pr.Kom.Mat. Przyr.Poznan.Tow.Przyj.Nauk.*, (1962), 10, 15.

- 35 In process variant (vi) the reaction is carried out in the presence of a base, preferably organometallic or metal hydride e.g. NaH, lithium diisopropylamide or NaOEt,

preferably in an aprotic solvent, preferably THF, ether or benzene at -78 to 25°C (analogous process in Gutswiller et al. (1978) J. Am. Chem. Soc. 100, 576).

In process variants (vii) and (viii) if a base is used it is preferably NaH, KH, an alkyl lithium e.g. BuLi, a metal alkoxide e.g. NaOEt, sodamide or lithium dialkylamide e.g. di-isopropylamide. An analogous method is described in US 3989691 and M. Gates et. al. (1970) J. Amer. Chem. Soc., 92, 205, as well as Taylor et al. (1972) JACS 94, 6218.

In process variant (x) where X or Y is CHO the reaction is a standard reductive alkylation using, e.g., sodium borohydride or sodium triacetoxyborohydride (Gribble, G. W. in *Encyclopedia of Reagents for Organic Synthesis* (Ed. Paquette, L. A.) (John Wiley and Sons, 1995), p 4649).

The process variants (xi) and (xii) are standard alkylation reactions well known to those skilled in the art, for example where an alcohol or amine is treated with an alkyl halide in the presence of a base (for example see March, J; *Advanced Organic Chemistry, Edition 3* (John Wiley and Sons, 1985), p364-366 and p342-343). The process is preferably carried out in a polar solvent such as N,N-dimethylformamide

In process variant (xiii) the reaction is a standard sulphonamide formation reaction well known to those skilled in the art. This may be e.g. the reaction of a sulphonyl halide with an amine.

In process variant (xiv) where X is W such as halogen, methanesulphonyloxy or trifluoromethanesulphonyloxy, the hydroxy group in Y is preferably converted to an OM group where M is an alkali metal by treatment of an alcohol with a base. The base is preferably inorganic such as NaH, lithium diisopropylamide or sodium. Where X is OH, the hydroxy group in Y is activated under Mitsunobu conditions (Fletcher et.al. J Chem Soc. (1995), 623). Alternatively the X=O and Y=CH₂OH groups can be reacted directly by activation with dichlorocarbodiimide (DCC) (Chem. Berichte 1962, 95, 2997 or Angewante Chemie 1963 75, 377).

In process variant (xv) the reaction is conducted in the presence of an organic base such as triethylamine or pyridine such as described by Fuhrman et.al., J. Amer. Chem. Soc.; 67, 1245, 1945. The X=NR^{11'}SO₂W or Y=SO₂W intermediates can be formed from the requisite amine e.g. by reaction with SO₂Cl₂ analogously to the procedure described by the same authors Fuhrman et.al., J. Amer. Chem. Soc.; 67, 1245, 1945.

In process variant (xvi) the reaction is an alkylation, examples of which are described in J. Med. chem. (1979) 22(10) 1171-6. The compound of formula (IV) may be prepared from the corresponding compound where X is NHR^{11'} by acylation with an appropriate derivative of the acid WCH₂COOH such as the acid chloride or sulphonation

with an appropriate derivative of the sulphonic acid WCH_2SO_3H such as the sulphonyl chloride.

In process variant (xvii) the leaving group W is preferably chloro or trifluoromethylsulphonyl and the reaction is the palladium catalysed process known as the "Buchwald" reaction (J. Yin and S. L. Buchwald, *Org.Lett.*, 2000, 2, 1101).

Reduction of a carbonyl group A or B to $CHOH$ can be readily accomplished using reducing agents well known to those skilled in the art, e.g. sodium borohydride in aqueous ethanol or lithium aluminium hydride in ethereal solution. This is analogous to methods described in EP53964, US384556 and J. Gutzwiller *et al*, *J. Amer. Chem. Soc.*, 1978, 100, 576.

The carbonyl group A or B may be reduced to CH_2 by treatment with a reducing agent such as hydrazine in ethylene glycol, at e.g. 130-160°C, in the presence of potassium hydroxide.

Reaction of a carbonyl group A or B with an organometallic reagent yields a group where R^8 is OH and R^9 is alkyl.

A hydroxy group on A or B may be oxidised to a carbonyl group by oxidants well known to those skilled in the art, for example, manganese dioxide, pyridinium chlorochromate or pyridinium dichromate.

A hydroxyalkyl A-B group CHR^7CR^9OH or $CR^7(OH)CHR^9$ may be dehydrated to give the group $CR^7=CR^9$ by treatment with an acid anhydride such as acetic anhydride.


Methods for conversion of $CR^7=CR^9$ by reduction to CHR^7CHR^9 are well known to those skilled in the art, for example using hydrogenation over palladium on carbon as catalyst. Methods for conversion of $CR^7=CR^9$ to give the A-B group $CR^7(OH)CHR^9$ or CHR^7CR^9OH are well known to those skilled in the art for example by epoxidation and subsequent reduction by metal hydrides, hydration, hydroboration or oxymercuration.

An amide carbonyl group may be reduced to the corresponding amine using a reducing agent such as lithium aluminium hydride.

A hydroxy group in A or B may be converted to azido by activation and displacement e.g. under Mitsunobu conditions using hydrazoic acid or by treatment with diphenylphosphorylazide and base, and the azido group in turn may be reduced to amino by hydrogenation.

An example of a group Q^1 convertible to NR^2R^4 is NR^2R^4 or halogen. Halogen may be displaced by an amine HNR^2R^4 by a conventional alkylation.

When Q^1 Q^2 together form a protected oxo group this may be an acetal such as ethylenedioxy which can subsequently be removed by acid treatment to give a compound of formula (VI):



wherein the variables are as described for formula (I)

Intermediates of formula (VI) are novel and as such form part of the invention.

The ketone of formula (VI) is reacted with an amine HNR^2R^4 by conventional reductive alkylation as described above for process variant (x).

Examples of groups Z^1 , Z^2 , Z^3 , Z^4 , Z^5 convertible to Z^1 , Z^2 , Z^3 , Z^4 and Z^5 include $\text{CR}^{1a'}$ where $\text{R}^{1a'}$ is a group convertible to R^{1a} . Z^1 , Z^2 , Z^3 , Z^4 and Z^5 are preferably Z^1 , Z^2 , Z^3 , Z^4 and Z^5 .

$\text{R}^{1a'}$, $\text{R}^{1'}$ and $\text{R}^{2'}$ are preferably R^{1a} , R^1 and R^2 . $\text{R}^{1'}$ is preferably methoxy. $\text{R}^{2'}$ is preferably hydrogen. $\text{R}^{3'}$ is R^3 or more preferably hydrogen, vinyl, alkoxy carbonyl or carboxy. $\text{R}^{4'}$ is R^4 or more preferably H or an N-protecting group such as t-butoxy carbonyl, benzyloxy carbonyl or 9-fluorenylmethyloxycarbonyl.

Conversions of $\text{R}^{1'}$, $\text{R}^{2'}$, $\text{R}^{3'}$ and $\text{R}^{4'}$ and interconversions of R^1 , R^2 , R^3 and R^4 are conventional. In compounds which contain an optionally protected hydroxy group, suitable conventional hydroxy protecting groups which may be removed without disrupting the remainder of the molecule include acyl and alkylsilyl groups. N-protecting groups are removed by conventional methods.

For example $\text{R}^{1'}$ methoxy is convertible to $\text{R}^{1'}$ hydroxy by treatment with lithium and diphenylphosphine (general method described in Ireland *et al*, *J. Amer. Chem. Soc.*, 1973, 7829) or HBr. Alkylation of the hydroxy group with a suitable alkyl derivative bearing a leaving group such as halide and a protected amino, piperidyl, amidino or guanidino group or group convertible thereto, yields, after conversion/deprotection, R^1 alkoxy substituted by optionally N-substituted amino, piperidyl, guanidino or amidino.

Substituted 2-oxo-oxazolidinyl containing R^3 groups may be prepared from the corresponding aldehyde by conventional reaction with a glycine anion equivalent, followed by cyclisation of the resulting amino alcohol (M. Grauert *et al*, *Ann. Chem.*, 1985, 1817; Rozenberg *et al*, *Angew. Chem. Int. Ed. Engl.*, 1994, 33(1), 91). The resulting 2-oxo-oxazolidinyl group contains a carboxy group which can be converted to other R^{10} groups by standard procedures.

Carboxy groups within R^3 may be prepared by Jones' oxidation of the corresponding alcohols CH_2OH using chromium acid and sulphuric acid in water/methanol (E.R.H. Jones *et al*, *J. Chem. Soc.*, 1946, 39). Other oxidising agents may be used for this transformation such as sodium periodate catalysed by ruthenium

trichloride (G.F. Tutwiler *et al*, *J. Med. Chem.*, 1987, 30(6), 1094), chromium trioxide-pyridine (G. Just *et al*, *Synth. Commun.*, 1979, 9(7), 613), potassium permanganate (D.E. Reedich *et al*, *J. Org. Chem.*, 1985, 50(19), 3535), and pyridinium chlorochromate (D. Askin *et al*, *Tetrahedron Lett.*, 1988, 29(3), 277).

5 The carboxy group may alternatively be formed in a two stage process, with an initial oxidation of the alcohol to the corresponding aldehyde using for instance dimethyl sulphoxide activated with oxalyl chloride (N.Cohen *et al*, *J. Am. Chem. Soc.*, 1983, 105, 3661) or dicyclohexylcarbodiimide (R.M.Wengler, *Angew. Chim. Int. Ed. Eng.*, 1985, 24(2), 77), or oxidation with tetrapropylammonium perruthenate (Ley *et al*, *J. Chem.Soc. Chem Commun.*, 1987, 1625). The aldehyde may then be separately oxidised to the

10 corresponding acid using oxidising agents such as silver (II) oxide (R.Grigg *et al*, *J. Chem. Soc. Perkin1*, 1983, 1929), potassium permanganate (A.Zurcher, *Helv. Chim. Acta.*, 1987, 70 (7), 1937), sodium periodate catalysed by ruthenium trichloride (T.Sakata *et al*, *Bull. Chem. Soc. Jpn.*, 1988, 61(6), 2025), pyridinium chlorochromate (R.S.Reddy *et al*, *Synth. Commun.*, 1988, 18(51), 545) or chromium trioxide (R.M.Coates *et al*, *J. Am. Chem. Soc.*, 1982, 104, 2198).

15 An R³ CO₂H group may also be prepared from oxidative cleavage of the corresponding diol, CH(OH)CH₂OH, using sodium periodate catalysed by ruthenium trichloride with an acetonitrile-carbontetrachloride-water solvent system (V.S.Martin *et al*, *Tetrahedron Letters*, 1988, 29(22), 2701).

20 Other routes to the synthesis of carboxy groups within R³ are well known to those skilled in the art.

R³ groups containing a cyano or carboxy group may be prepared by conversion of an alcohol to a suitable leaving group such as the corresponding tosylate by reaction with 25 para-toluenesulphonyl chloride (M.R. Bell, *J. Med. Chem.*, 1970, 13, 389), or the iodide using triphenylphosphine, iodine, and imidazole (G. Lange, *Synth. Commun.*, 1990, 20, 1473). The second stage is the displacement of the leaving group with cyanide anion (L.A. Paquette *et al*, *J. Org. Chem.*, 1979, 44(25), 4603; P.A. Grieco *et al*, *J. Org. Chem.*, 1988, 53(16), 3658. Finally acidic hydrolysis of the nitrile group gives the desired acids 30 (H.Rosemeyer *et al*, *Heterocycles*, 1985, 23 (10), 2669). The hydrolysis may also be carried out with base e.g. potassium hydroxide (H.Rapoport, *J. Org. Chem.*, 1958, 23, 248) or enzymatically (T. Beard *et al*, *Tetrahedron Asymmetry*, 1993, 4 (6), 1085).


R^{3'} cis or trans hydroxy may be introduced by the methods of van Deale *et al.*, *Drug Development Research* 8:225-232 (1986) or *Heterocycles* 39(1), 163-170 (1994).

35 For trans hydroxy, a suitable method converts N-protected tetrahydropyridine to the epoxide by treatment with metachloroperbenzoic acid, followed by opening of the epoxide with a suitable amine NR^{2'}R^{4'}. R^{3'} hydroxy may then be converted to optionally

substituted amino via preparation of the R^{3'} amino derivative by standard transformations such as a Mitsunobu reaction (for example as reviewed in Misunobu, Synthesis, (1981), 1), for example with succinimide in the presence of diethylazodicarboxylate and triphenylphosphine to give the phthalimidoethylpiperidine. Removal of the phthaloyl group, for example by treatment with methylhydrazine, affords the R^{3'} amine. Optional substitution may then be introduced by standard methods for amine substitution well known to those skilled in the art.

R³ 4-CF₃ may be introduced by the following scheme I:

Scheme I




(a) (Boc)₂O, CH₂Cl₂; (b) LDA, then CF₃-X; (c) NaOH, H₂O, EtOH; (d) DPPA, Et₃N, toluene, then BnOH; (e) H₂, Pd/C, EtOH.

Commercially-available ethyl isonipecotate (I-1) reacts with an appropriate acylating agent, preferably di-*tert*-butyl dicarbonate, to afford the protected derivative I-2. Typical solvents for this reaction include CH₂Cl₂, THF, or DMF. The protecting group for the amine must be compatible with subsequent chemistry, and must be readily removable when desired. Methods for the protection of amines are well-known to those of skill in the art, and are described in standard reference volumes, such as Greene "Protective Groups in Organic Synthesis" (published by Wiley-Interscience). Alkylation of I-2 can be accomplished by reaction with an appropriate base, typically LDA or LiN(TMS)₂, in an aprotic solvent, usually THF or DME, followed by trapping of the enolate with an appropriate electrophile, to afford I-3. Trifluoromethyl iodide (CF₃I) or *S*-(trifluoromethyl)dibenzothiophenium trifluoromethanesulfonate are typically preferred as electrophilic trifluoromethylating reagents. The ethyl ester of I-3 is hydrolyzed using aqueous base, for example, LiOH in aqueous THF or NaOH in aqueous methanol or ethanol, and the intermediate carboxylate salt is acidified with a suitable acid, for instance TFA or HCl, to afford the carboxylic acid I-4. Curtius-type rearrangement of I-4 gives an intermediate isocyanate, which typically is not isolated, but rather is reacted *in situ* with

an appropriate alcohol, such as benzyl alcohol, to give **I-5**. Diphenylphosphoryl azide in the presence of an amine base, generally triethylamine or diisopropylethylamine (Hunig's base), is the preferred reagent combination for effecting the Curtius-type rearrangement of **I-4**, but more classical conditions, such as formation of the acid chloride, reaction with azide anion, and warming of the acyl azide, can also be used. The benzyloxycarbonyl group in **I-5** is removed by hydrogenolysis in the presence of a palladium catalyst, typically palladium on activated charcoal, in a suitable solvent, usually EtOH, MeOH, EtOAc, or mixtures thereof, to give amine **I-6**.

R^3 2-CF₃ may be introduced by the following scheme II:

Scheme II




- 15 (a) 1-methoxy-3-(trimethylsilyloxy)-1,3-butadiene, ZnCl₂, CH₃CN; (b) separate diastereomers; (c) L-Selectride®, THF; (d) NH₂OCH₃, *p*-TsOH, toluene; (e) LiAlH₄, THF; (f) (Boc)₂O, CH₂Cl₂; (g) H₂, Pd/C, EtOH.

Imine **II-1**, prepared in standard fashion by acid-catalyzed reaction of trifluoroacetaldehyde ethyl hemiacetal and (*R*)-(+)- α -methylbenzylamine, reacts with a silyloxydiene, for example 1-methoxy-3-(trimethylsilyloxy)-1,3-butadiene, in a Diels-Alder reaction to afford piperidone **II-2**. The reaction is conducted in a neutral solvent such as CH₃CN, THF, or CH₂Cl₂, and oftentimes is mediated by a Lewis acid such as ZnCl₂. Diastereomers are best separated at this point. The enone **II-2** is reduced to the

corresponding ketone **II-3** by reaction with L-Selectride® in a suitable solvent, generally THF or DME, and the ketone is converted to an oxime derivative under standard conditions well-known to those of skill in the art. Reduction of the oxime derivative under standard conditions (LiAlH_4) gives a mixture of diastereomeric amines from which the amine **II-5** can be isolated. The amine is protected with an appropriate protecting group, preferably a *tert*-butyl carbamate (see Scheme I), to afford **II-6**. The α -methylbenzyl group of **II-6** is removed by hydrogenolysis in the presence of a palladium catalyst, typically palladium on activated charcoal, in a suitable solvent, usually EtOH, MeOH, EtOAc, or mixtures thereof, to give amine **II-7**.

R^3 3- CF_3 may be introduced by the following scheme III:

Scheme III



(a) TMSCl , Et_3N , DMF; (b) $\text{CF}_3\text{-X}$, DMF; (c) (*R*)-(+)- α -methylbenzylamine, *p*-TsOH, toluene; (d) NaBH_4 , EtOH; (e) separate diastereomers; (f) H_2 , Pd/C, EtOH.


The commercially-available ketone **III-1** is converted to the corresponding silyl enol ether **III-2** by reaction with a silylating reagent, such as trimethylsilyl chloride or trimethylsilyl triflate, in the presence of an amine base, typically triethylamine, in a suitable solvent, such as diethyl ether, THF, DMF, or mixtures thereof. The silyl enol ether **III-2** reacts with an electrophilic trifluoromethylating reagent, such as trifluoromethyl iodide (CF_3I) or more preferably *S*-(trifluoromethyl)dibenzothiophenium trifluoromethanesulfonate (see *Tet. Lett.* 1990, 31, 3579-3582)), in an appropriate solvent, such as THF, DMF, or mixtures thereof, to afford the α -trifluoromethyl ketone **III-3**. Ketone **III-3** reacts with a chiral amine, for instance (*R*)-(+)- α -methylbenzylamine, under standard acidic catalysis, to afford the imine derivative **III-4**, which can be reduced to afford amine **III-5**. This type of reduction is typically conducted using sodium borohydride, sodium cyanoborohydride or sodium (triacetoxo)borohydride, in an appropriate solvent, such as EtOH, MeOH, THF, CH_2Cl_2 , $\text{ClCH}_2\text{CH}_2\text{Cl}$, or mixtures

thereof. Diastereomers are best separated at this point. The α -methylbenzyl group of **III-5** is removed by hydrogenolysis in the presence of a palladium catalyst, typically palladium on activated charcoal, in a suitable solvent, usually EtOH, MeOH, EtOAc, or mixtures thereof, to give amine **III-6**.

5

R^3 2-oxo may be introduced by the following scheme IV:

Scheme IV



10


(a) NaH, THF, 0 °C to RT; (b) 10% Pd/C, H₂, MeOH.

(*R,S*)-4-(Dibenzylamino)piperidin-2-one (**IV-2**, Homo-Freidinger Lactam, prepared from (*R,S*)-aspartic acid according to the procedure of Weber and Gmeiner, *Synlett*, 1998, 885-887) reacts with an appropriate epoxide, for instance 6-methoxy-4-(*R*)-oxiranylquinoline (**VI-1**) or 6-methoxy-4-(*R*)-oxiranyl-[1,5]naphthyridine, to afford the adduct **IV-3**. The reaction is mediated by a strong base, preferably sodium hydride, which is used to deprotonate **IV-2**, and is typically conducted in a polar, aprotic solvent, such as THF, DMF, or mixtures thereof. The benzyl groups in **IV-3** are removed by hydrogenolysis in the presence of a palladium catalyst, typically palladium on activated charcoal, in a suitable solvent, usually EtOH, MeOH, EtOAc, or mixtures thereof, to give amine **IV-4**.

20

R^3 3-F may be introduced by the following scheme V:

Scheme V



- 5 (a) TMSCl, Et₃N, DMF, 80 °C; (b) Selectfluor, CH₃CN; (c) benzylamine, 1,2-dichloroethane, Na(OAc)₃BH; (d) separate diastereomers; (e) 10% Pd/C, H₂, HCl, EtOH.

The trimethylsilyl enol ether (V-2), prepared from commercially-available *N*-(*tert*-butoxycarbonyl)piperidone (V-1) as described in Scheme III, reacts with an electrophilic
 10 fluorinating reagent, preferably Selectfluor (1-chloromethyl-4-fluoro-1,4-diazabicyclo[2.2.2]octane bis(tetrafluoroborate), in a neutral solvent such as CH₃CN, to afford the α -fluoro ketone V-3. Reductive amination of V-3 with benzylamine according to the procedures described in Schemes I and III gives the expected 4-aminobenzyl-3-
 15 fluoro-*N*-(*tert*-butoxycarbonyl)piperidine derivatives V-4 and V-5 as a mixture of *cis*- and *trans*-isomers in an 8:1 ratio. These diastereomers are separable by chromatography on silica gel. The predominate *cis*-mixture of enantiomers is debenzylated by catalytic hydrogenation as described in Scheme II, to give the amino derivative V-6.

Other functional groups in R³ may be obtained by conventional conversions of hydroxy, carboxy or cyano groups.

20 Tetrazoles are conveniently prepared by reaction of sodium azide with the cyano group (e.g. F. Thomas *et al*, *Bioorg. Med. Chem. Lett.*, 1996, 6(6), 631; K. Kubo *et al*, *J. Med. Chem.*, 1993, 36, 2182) or by reaction of azidotri-*n*-butyl stannane with the cyano group followed by acidic hydrolysis (P.L. Ornstein, *J. Org. Chem.*, 1994, 59, 7682 and *J. Med. Chem.*, 1996, 39 (11), 2219).

25 The 3-hydroxy-3-cyclobutene-1,2-dion-4-yl group (e.g. R.M. Soll, *Bioorg. Med. Chem. Lett.*, 1993, 3(4), 757 and W.A. Kinney, *J. Med. Chem.*, 1992, 35(25), 4720) can be prepared by the following sequence:- (1) a group (CH₂)_nCHO (n = 0,1,2) is treated with triethylamine, carbon tetrabromide-triphenylphosphine to give initially (CH₂)_nCH=CHBr; (2) dehydrobromination of this intermediate to give the corresponding
 30 bromoethyne derivative (CH₂)_nC \equiv CBr (for this 2 stage sequence see D. Grandjean *et al*, *Tetrahedron Lett.*, 1994, 35(21), 3529); (3) palladium-catalysed coupling of the

bromoethyne with 4-(1-methylethoxy)-3-(tri-n-butylstannyl)cyclobut-3-ene-1,2-dione (Liebeskind *et al*, *J. Org. Chem.*, 1990, 55, 5359); (4) reduction of the ethyne moiety to -CH₂CH₂- under standard conditions of hydrogen and palladium on charcoal catalysis (see Howard *et al*, *Tetrahedron*, 1980, 36, 171); and finally (4) acidic hydrolysis of the methyl ethoxyester to generate the corresponding 3-hydroxy-3-cyclobutene-1,2-dione group (R.M. Soll, *Bioorg. Med. Chem. Lett.*, 1993, 3(4), 757).

The tetrazol-5-ylaminocarbonyl group may be prepared from the corresponding carboxylic acid and 2-aminotetrazole by dehydration with standard peptide coupling agents such as 1,1'-carbonyldiimidazole (P.L. Ornstein *et al*, *J. Med Chem*, 1996, 39(11), 2232).

The alkyl- and alkenyl-sulphonylcarboxamides are similarly prepared from the corresponding carboxylic acid and the alkyl- or alkenyl-sulphonamide by dehydration with standard peptide coupling agents such as 1,1'-carbonyldiimidazole (P.L. Ornstein *et al*, *J. Med. Chem.*, 1996, 39(11), 2232).

The hydroxamic acid groups are prepared from the corresponding acids by standard amide coupling reactions e.g. N.R. Patel *et al*, *Tetrahedron*, 1987, 43(22), 5375.

2,4-Thiazolidinedione groups may be prepared from the aldehydes by condensation with 2,4-thiazolidinedione and subsequent removal of the olefinic double bond by hydrogenation.

The preparation of 5-oxo-1,2,4-oxadiazoles from nitriles is described by Y. Kohara *et al*, *Bioorg. Med. Chem. Lett.*, 1995, 5(17), 1903.

1,2,4-Triazol-5-yl groups may be prepared from the corresponding nitrile by reaction with an alcohol under acid conditions followed by reaction with hydrazine and then an R¹⁰-substituted activated carboxylic acid (see J.B. Polya in "Comprehensive Heterocyclic Chemistry" Edition 1, p762, Ed A.R. Katritzky and C.W. Rees, Pergamon Press, Oxford, 1984 and J.J. Ares *et al*, *J. Heterocyclic Chem.*, 1991, 28(5), 1197).

Other substituents on R³ alkyl or alkenyl may be interconverted by conventional methods, for example hydroxy may be derivatised by esterification, acylation or etherification. Hydroxy groups may be converted to halogen, thiol, alkylthio, azido, alkylcarbonyl, amino, aminocarbonyl, oxo, alkylsulphonyl, alkenylsulphonyl or aminosulphonyl by conversion to a leaving group and substitution by the required group or oxidation as appropriate or reaction with an activated acid, isocyanate or alkoxyisocyanate. Primary and secondary hydroxy groups can be oxidised to an aldehyde or ketone respectively and alkylated with a suitable agent such as an organometallic reagent to give a secondary or tertiary alcohol as appropriate. A carboxylate group may be converted to a hydroxymethyl group by reduction of an ester of this acid with a suitable reducing agent such as LiAlH₄.

An NH_2 substituent on piperidine is converted to NR^2R^4 by conventional means such as amide or sulphonamide formation with an acyl derivative R^5COW or $\text{R}^5\text{SO}_2\text{W}$, for compounds where U is CO or SO_2 or, where U is CH_2 , alkylation with an alkyl halide R^5CH_2 -halide in the presence of base, acylation/reduction with an acyl derivative R^5COW or reductive alkylation with an aldehyde R^5CHO .

Where one of R^3 and R^6 , R^7 , R^8 or R^9 contains a carboxy group and the other contains a hydroxy or amino group they may together form a cyclic ester or amide linkage. This linkage may form spontaneously during coupling of the compound of formula (IV) and the piperidine moiety or in the presence of standard peptide coupling agents.

It will be appreciated that under certain circumstances interconversions may interfere, for example, A or B hydroxy groups in A or B and the piperidine substituent NH_2 will require protection e.g. as a carboxy- or silyl-ester group for hydroxy and as an acyl derivative for piperidine NH_2 , during conversion of R^1 , R^2 , R^3 or R^4 , or during the coupling of the compounds of formulae (IV) and (V).

Compounds of formulae (IV) and (V) are known compounds, (see for example Smith *et al*, *J. Amer. Chem. Soc.*, 1946, 68, 1301) or prepared analogously.

Compounds of formula (IV) where X is $\text{CR}^6\text{R}^7\text{SO}_2\text{W}$ may be prepared by a route analogous to that of Ahmed El Hadri *et al*, *J. Heterocyclic Chem.*, 1993, 30(3), 631. Thus compounds of formula (IV) where X is $\text{CH}_2\text{SO}_2\text{OH}$ may be prepared by reacting the corresponding 4-methyl compound with N-bromosuccinimide, followed by treatment with sodium sulfite. The leaving group W may be converted to another leaving group W, e.g. a halogen group, by conventional methods.

The isocyanate of formula (IV) may be prepared conventionally from a 4-amino derivative such as 4-amino-quinoline, and phosgene, or phosgene equivalent (eg triphosgene) or it may be prepared more conveniently from a 4-carboxylic acid by a "one-pot" Curtius Reaction with diphenyl phosphoryl azide (DPPA) [see T. Shiori *et al*. *Chem. Pharm. Bull.* 35, 2698-2704 (1987)].

The 4-amino derivatives are commercially available or may be prepared by conventional procedures from a corresponding 4-chloro or 4-trifluoromethanesulphonate derivative by treatment with ammonia (O.G. Backeberg *et al.*, *J. Chem Soc.*, 381, 1942) or propylamine hydrochloride (R. Radinov *et al.*, *Synthesis*, 886, 1986).

4-Alkenyl compounds of formula (IV) may be prepared by conventional procedures from a corresponding 4-halogeno-derivative by e.g. a Heck-synthesis as

described in e.g. *Organic Reactions*, 1982, 27, 345.

4-Halogeno derivatives of compounds of formula (IV) are commercially available, or may be prepared by methods known to those skilled in the art. A 4-chloroquinoline is

prepared from the corresponding quinolin-4-one by reaction with phosphorus oxychloride (POCl_3) or phosphorus pentachloride, PCl_5 . A 4-chloroquinazoline is prepared from the corresponding quinazolin-4-one by reaction with phosphorus oxychloride (POCl_3) or phosphorus pentachloride, PCl_5 . A quinazolinone and quinazolines may be prepared by standard routes as described by T.A. Williamson in *Heterocyclic Compounds*, 6, 324 (1957) Ed. R.C. Elderfield.

Activated carboxy derivatives $\text{X}=\text{A}'\text{COW}$ of formula (IV) may be prepared from $\text{X}=\text{A}'\text{CO}_2\text{H}$ derivatives in turn prepared from CO_2H derivatives by conventional methods such as homologation.

10 ~~4-Carboxy derivatives of compounds of formula (IV) are commercially available~~
or may be prepared by conventional procedures for preparation of carboxy heteroaromatics well known to those skilled in the art. For example, quinazolines may be prepared by standard routes as described by T.A. Williamson in *Heterocyclic Compounds*, 6, 324 (1957) Ed. R.C. Elderfield. These 4-carboxy derivatives may be
15 activated by conventional means, e.g. by conversion to an acyl halide or anhydride.

Pyridazines may be prepared by routes analogous to those described in Comprehensive Heterocyclic Chemistry, Volume 3, Ed A.J. Boulton and A. McKillop and naphthyridines may be prepared by routes analogous to those described in Comprehensive Heterocyclic Chemistry, Volume 2, Ed A.J. Boulton and A. McKillop.
20 A 4-oxirane derivative of compounds of formula (IV) is conveniently prepared from the 4-carboxylic acid by first conversion to the acid chloride with oxalyl chloride and then reaction with trimethylsilyldiazomethane to give the diazoketone derivative. Subsequent reaction with 5M hydrochloric acid gives the chloromethylketone. Reduction with sodium borohydride in aqueous methanol gives the chlorohydrin which undergoes
25 ring closure to afford the epoxide on treatment with base, e.g. potassium hydroxide in ethanol-tetrahydrofuran.

Alternatively and preferably, 4-oxirane derivatives can be prepared from bromomethyl ketones which can be obtained from 4-hydroxy compounds by other routes well known to those skilled in the art. For example, hydroxy compounds can be converted
30 to the corresponding 4-trifluoromethanesulphonates by reaction with trifluoromethanesulphonic anhydride under standard conditions (see K. Ritter, *Synthesis*, 1993, 735). Conversion into the corresponding butyloxyvinyl ethers can be achieved by a Heck reaction with butyl vinyl ether under palladium catalysis according to the procedure of W. Cabri *et al*, *J. Org. Chem*, 1992, 57 (5), 1481. (Alternatively, the same
35 intermediates can be attained by Stille coupling of the trifluoromethanesulphonates or the analogous chloro derivatives with (1-ethoxyvinyl)tributyl tin, T. R. Kelly, *J. Org. Chem.*, 1996, 61, 4623.) The alkyloxyvinyl ethers are then converted into the corresponding

bromomethylketones by treatment with N-bromosuccinimide in aqueous tetrahydrofuran in a similar manner to the procedures of J. F. W. Keana, *J. Org. Chem.*, 1983, **48**, 3621 and T. R. Kelly, *J. Org. Chem.*, 1996, **61**, 4623.

5 The 4-hydroxyderivatives can be prepared from an aminoaromatic by reaction with methylpropiolate and subsequent cyclisation, analogous to the method described in N. E. Heindel et al, *J. Het. Chem.*, 1969, **6**, 77. For example, 5-amino-2-methoxy pyridine can be converted to 4-hydroxy-6-methoxy-[1,5]naphthyridine using this method.

10 If a chiral reducing agent such as (+) or (-)-B-chlorodiisopinocampheylborane ['DIP-chloride'] is substituted for sodium borohydride, the prochiral chloromethylketone is converted into the chiral chlorohydrin with ee values generally 85-95% [see C. Bolm *et al*, *Chem. Ber.* **125**, 1169-1190, (1992)]. Recrystallisation of the chiral epoxide gives material in the mother liquor with enhanced optical purity (typically ee 95%).

The (*R*)-epoxide, when reacted with a piperidine derivative gives ethanolamine compounds as single diastereomers with (*R*)-stereochemistry at the benzylic position.

15 Alternatively, the epoxide may be prepared from the 4-carboxaldehyde by a Wittig approach using trimethylsulfonium iodide [see G.A. Epling and K-Y Lin, *J. Het. Chem.*, 1987, **24**, 853-857], or by epoxidation of a 4-vinyl derivative.

20 4-Hydroxy-1,5-naphthyridines can be prepared from 3-aminopyridine derivatives by reaction with diethyl ethoxymethylene malonate to produce the 4-hydroxy-3-carboxylic acid ester derivative with subsequent hydrolysis to the acid, followed by thermal decarboxylation in quinoline (as for example described for 4-Hydroxy-[1,5]naphthyridine-3-carboxylic acid, J. T. Adams *et al.*, *J. Amer. Chem. Soc.*, 1946, **68**, 1317). A 4-hydroxy-[1,5]naphthyridine can be converted to the 4-chloro derivative by heating in phosphorus oxychloride, or to the 4-methanesulphonyloxy or 4-
25 trifluoromethanesulphonyloxy derivative by reaction with methanesulphonyl chloride or trifluoromethanesulphonic anhydride, respectively, in the presence of an organic base. A 4-amino 1,5-naphthyridine can be obtained from the 4-chloro derivative by reaction with n-propylamine in pyridine.

30 Similarly, 6-methoxy-1,5-naphthyridine derivatives can be prepared from 3-amino-6-methoxypyridine.

1,5-Naphthyridines may be prepared by other methods well known to those skilled in the art (for examples see P.A. Lowe in "Comprehensive Heterocyclic Chemistry" Volume 2, p581-627, Ed A.R. Katritzky and C.W. Rees, Pergamon Press, Oxford, 1984).

35 The 4-hydroxy and 4-amino-cinnolines may be prepared following methods well known to those skilled in the art [see A.R. Osborn and K. Schofield, *J. Chem. Soc.* 2100 (1955)]. For example, a 2-aminoacetophenone is diazotised with sodium nitrite and acid

to produce the 4-hydroxycinnoline with conversion to chloro and amino derivatives as described for 1,5-naphthyridines.

For compounds of formula (V), suitable amines may be prepared from the corresponding 4-substituted piperidine acid or alcohol. In a first instance, an N-protected piperidine containing an acid bearing substituent, can undergo a Curtius rearrangement and the intermediate isocyanate can be converted to a carbamate by reaction with an alcohol. Conversion to the amine may be achieved by standard methods well known to those skilled in the art used for amine protecting group removal. For example, an acid substituted N-protected piperidine can undergo a Curtius rearrangement e.g. on treatment with diphenylphosphoryl azide and heating, and the intermediate isocyanate reacts in the presence of 2-trimethylsilylethanol to give the trimethylsilylethylcarbamate (T.L. Capson & C.D. Poulter, *Tetrahedron Lett.*, 1984, 25, 3515). This undergoes cleavage on treatment with tetrabutylammonium fluoride to give the 4-amine substituted N-protected piperidine.

In a second instance, an N-protected piperidine containing an alcohol bearing substituent undergoes a Mitsunobu reaction (for example as reviewed in Mitsunobu, *Synthesis*, (1981), 1), for example with succinimide in the presence of diethyl azodicarboxylate and triphenylphosphine to give the phthalimidoethylpiperidine. Removal of the phthaloyl group, for example by treatment with methylhydrazine, gives the amine of formula (V).

R^5CH_2 -halides, acyl derivative R^5COW and R^5SO_2W or aldehydes R^5CHO are commercially available or are prepared conventionally. The aldehydes may be prepared by partial reduction of the R^5 -ester with lithium aluminium hydride or diisobutylaluminium hydride or more preferably by reduction to the alcohol, with lithium aluminium hydride or sodium borohydride, followed by oxidation to the aldehyde with manganese (II) dioxide. The aldehydes may also be prepared from carboxylic acids in two stages by conversion to a mixed anhydride for example by reaction with isobutyl chloroformate followed by reduction with sodium borohydride (R. J. Alabaster et al., *Synthesis*, 598, 1989) to give the hydroxymethyl substituted heteroaromatic or aromatic and then oxidation with a standard oxidising agent such as pyridinium dichromate or manganese (II) dioxide. Acyl derivative R^5COW may be prepared by activation of the R^5 -acid. R^5CH_2 -halides such as bromides may be prepared from the alcohol R^5CH_2OH by reaction with phosphorus tribromide in DCM/triethylamine. Alternatively the aldehyde R^5CHO and sulphonic acid derivative R^5SO_2W may be generated by treatment of the R^5H heterocycle with suitable reagents. For example by formylation with hexamine in either trifluoroacetic acid or methanesulfonic acid, in a modified Duff procedure [O. I. Petrov et al. *Collect. Czech. Chem. Commun.* 62, 494-497 (1997)]. Reaction of a R^5H

heterocycle with chlorosulphonic acid gives the sulphonic acid derivative (by methods analogous to Techer *et. al.*, *C.R.Hebd. Seances Acad. Sci. Ser.C*; 270, 1601, 1970)..

R⁵ heterocycles are commercially available or may be prepared by conventional methods.

5 The amines R²R⁴NH are available commercially or prepared conventionally. For example amines R⁵CH₂NH₂ may be prepared from a bromomethyl derivative by reaction with sodium azide in dimethylformamide (DMF), followed by hydrogenation of the azidomethyl derivative over palladium-carbon. An alternative method is to use potassium phthalimide/DMF to give the phthalimidomethyl derivative, followed by reaction with
10 hydrazine in DCM to liberate the primary amine.

Conversions of R^{1a'}, R^{1'}, R^{2'}, R^{3'} and R^{4'} may be carried out on the intermediates of formulae (IV), and (V) prior to their reaction to produce compounds of formula (I) in the same way as described above for conversions after their reaction.

15 Further details for the preparation of compounds of formula (I) are found in the examples.

The compounds of formula (I) may be prepared singly or as compound libraries comprising at least 2, for example 5 to 1,000 compounds, and more preferably 10 to 100 compounds of formula (I). Libraries of compounds of formula (I) may be prepared by a combinatorial "split and mix" approach or by multiple parallel synthesis using either
20 solution phase or solid phase chemistry, by procedures known to those skilled in the art.

Thus according to a further aspect of the invention there is provided a compound library comprising at least 2 compounds of formula (I) or pharmaceutically acceptable derivatives thereof.

Novel intermediates of formulae (IV) and (V) are also part of this invention.

25 The antibacterial compounds according to the invention may be formulated for administration in any convenient way for use in human or veterinary medicine, by analogy with other antibacterials.

The pharmaceutical compositions of the invention include those in a form adapted for oral, topical or parenteral use and may be used for the treatment of bacterial infection
30 in mammals including humans.

The composition may be formulated for administration by any route. The compositions may be in the form of tablets, capsules, powders, granules, lozenges, creams or liquid preparations, such as oral or sterile parenteral solutions or suspensions.

35 The topical formulations of the present invention may be presented as, for instance, ointments, creams or lotions, eye ointments and eye or ear drops, impregnated dressings and aerosols, and may contain appropriate conventional additives such as preservatives, solvents to assist drug penetration and emollients in ointments and creams.

The formulations may also contain compatible conventional carriers, such as cream or ointment bases and ethanol or oleyl alcohol for lotions. Such carriers may be present as from about 1% up to about 98% of the formulation. More usually they will form up to about 80% of the formulation.

5 Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinylpyrrolidone; fillers, for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricants, for
10 example magnesium stearate, talc, polyethylene glycol or silica; disintegrants, for example potato starch; or acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid
15 preparations may contain conventional additives, such as suspending agents, for example sorbitol, methyl cellulose, glucose syrup, gelatin, hydroxyethyl cellulose, carboxymethyl cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, oily esters such as glycerine, propylene
20 glycol, or ethyl alcohol; preservatives, for example methyl or propyl *p*-hydroxybenzoate or sorbic acid, and, if desired, conventional flavouring or colouring agents.

Suppositories will contain conventional suppository bases, e.g. cocoa-butter or other glyceride.

25 For parenteral administration, fluid unit dosage forms are prepared utilizing the compound and a sterile vehicle, water being preferred. The compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilised before filling into a suitable vial or ampoule and sealing.

30 Advantageously, agents such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. The dry lyophilized powder is then sealed in the vial and an accompanying vial of water for injection may be supplied to reconstitute the liquid prior to use. Parenteral suspensions are prepared in substantially the same manner except that the compound is suspended in
35 the vehicle instead of being dissolved and sterilization cannot be accomplished by filtration. The compound can be sterilised by exposure to ethylene oxide before

suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

5 The compositions may contain from 0.1% by weight, preferably from 10-60% by weight, of the active material, depending on the method of administration. Where the compositions comprise dosage units, each unit will preferably contain from 50-500 mg of the active ingredient. The dosage as employed for adult human treatment will preferably range from 100 to 3000 mg per day, for instance 1500 mg per day depending on the route and frequency of administration. Such a dosage corresponds to 1.5 to 50 mg/kg per day. Suitably the dosage is from 5 to 20 mg/kg per day.

10 No toxicological effects are indicated when a compound of formula (I) or a pharmaceutically acceptable derivative thereof is administered in the above-mentioned dosage range.

15 The compound of formula (I) may be the sole therapeutic agent in the compositions of the invention or a combination with other antibacterials. If the other antibacterial is a β -lactam then a β -lactamase inhibitor may also be employed.


Compounds of formula (I) are active against a wide range of organisms including both Gram-negative and Gram-positive organisms.

20 All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The following examples illustrate the preparation of certain compounds of formula (I) and the activity of certain compounds of formula (I) against various bacterial organisms.

EXAMPLES


Example 1. Preparation of 6-[(3*S*,4*R*)-3-fluoro-1-[(*R*)-2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]piperidin-4-ylamino)methyl]-[1,2,3]thiadiazolo[5,4-*b*]pyridine and 6-[(3*R*,4*S*)-3-fluoro-1-[(*R*)-2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]piperidin-4-ylamino)methyl]-[1,2,3]thiadiazolo[5,4-*b*]pyridine



(a) (3*R*,4*S*)- and (3*S*,4*R*)-4-Amino-1-*tert*-butoxycarbonyl-3-fluoropiperidine

To a solution of the enantiomeric mixture of *cis*-4-benzylamino-1-*tert*-butoxycarbonyl-3-fluoropiperidine (prepared according to the procedures of *J. Med. Chem.* 1999, 42, 2087-2104, 1.0 g, 3.2 mmole) in EtOH (40 mL) was added 3 N HCl (2.5 mL) and 10 % Pd/C (50 mg). The reaction was shaken under H₂ (40 psi) on a Parr apparatus for 14 h, then was filtered through celite®. The filtrate was concentrated under reduced pressure, and the residue was purified by flash chromatography on silica gel (10% MeOH/CHCl₃) to afford the title compound (370 mg, 53%) as a white solid: MS (ES) *m/e* 219 (M + H)⁺.

(b) [1,2,3]Thiadiazolo[5,4-*b*]pyridine-6-carboxaldehyde



(i) 5-Amino-6-thioxo-1,6-dihydro-pyridine-3-carboxylic acid methyl ester

A mixture of sodium sulfide nonahydrate (2.17g) and sulfur (0.29g) was heated in boiling water (20mL) until the solution was homogeneous and added to a solution of 6-chloro-5-nitro-nicotinic acid methyl ester (see Example 13a) (3.10g) in methanol (50mL). The mixture was boiled for 15 minutes and cooled. The resulting disulfide was collected and washed with water to give a yellow solid (2.46g). The solid (5g) in acetic acid (100mL) and 4M HCl in dioxan (50mL) was treated with zinc dust (12g) and the mixture was stirred at room temperature for 30 minutes, filtered and evaporated to dryness. Sodium acetate and sodium sulfate were added and the mixture was extracted with warm chloroform and chromatographed on silica gel, eluting with chloroform then methanol-chloroform to afford a yellow solid (2.3g).

MS (+ve ion electrospray) *m/z* 185(MH⁺)

(ii) [1,2,3]Thiadiazolo[5,4-*b*]pyridine-6-carboxylic acid methyl ester

The amine (i) (1.3 g) was suspended in 0.5 M hydrochloric acid (200mL) and cooled to -3°C. A solution of sodium nitrite (487 mg) in water (3 mL) was added dropwise over 10 minutes and the mixture was stirred for 2 hours when the solid product was collected and chromatographed on silica gel (chloroform) to afford a solid (0.90g)

5 MS (+ve ion electrospray) m/z 196 (MH⁺)

(iii) [1,2,3]Thiadiazolo[5,4-b]pyridine-6-carboxylic acid

The ester (ii) (0.94g) was hydrolysed with aqueous sodium hydroxide in tetrahydrofuran to afford a solid (0.84g).

MS (-ve ion electrospray) m/z 180 (M-H⁻).

10 (iv) [1,2,3]Thiadiazolo[5,4-b]pyridin-6-yl-methanol

The carboxylic acid (iii) (0.82g) was reacted with *isobutylchloroformate* and sodium borohydride to afford a semi-solid (0.12g), after chromatography on silica gel (chloroform).

(v) [1,2,3]Thiadiazolo[5,4-b]pyridine-6-carboxaldehyde

15 The alcohol (iv) (0.10 g) was oxidised with manganese dioxide to afford a solid (51 mg).

MS (+ve ion electrospray in methanol) m/z 198 (MH⁺ for methanol adduct)

20 (c) (3*R*,4*S*) and (3*S*,4*R*)-3-Fluoro-4-[[[1,2,3]thiadiazolo[5,4-b]pyridine-6-ylmethyl]-amino]-piperidine

A solution of *cis*-4-amino-1-*tert*-butoxycarbonyl-3-fluoropiperidine (1.00 mmole) in CH₂Cl₂ (5 mL) and EtOH (0.5 mL) is treated with anhydrous Na₂SO₄ (280 mg) and [1,2,3]thiadiazolo[5,4-b]pyridine-6-carboxaldehyde (1.10 mmole). The resulting solution is stirred at room temperature, then sodium triacetoxo borohydride (320 mg, 1.50 mmole) is added. The mixture is quenched by the addition of water (2 mL) and the volatiles removed *in vacuo*. Purification gives the desired compound.

(d) 6-Methoxyquinoline-4-carboxylic acid

30 The title compound was prepared by modification of the procedure described by W.E. Doering and J.D. Chanley, *J. Amer. Chem. Soc.*, 1946, 68, 586. A mixture of quinone (derived from quinine by reaction with potassium *tert*-butoxide and benzophenone in toluene) (225g, 0.70 mol), *tert*-butanol (1 litre) and water (10 ml) was treated with potassium *tert*-butoxide (170g, 1.5 mol). The mixture was stirred at 30°C, while air was bubbled through for 3 days. The mixture was diluted with diethyl ether and water and the layers separated. The aqueous phase was extracted with ethyl acetate. The combined diethyl ether and ethyl acetate extracts were dried over magnesium sulfate and evaporated to give recovered starting material (approximately 100g). The aqueous phase

was acidified to pH5 with 5M hydrochloric acid. The precipitate was collected by filtration, washed with water and methanol, then dried to give 6-methoxyquinoline-4-carboxylic acid as a yellow solid (64.6g, 46%).

5 δ H (d-6 DMSO) 6.23-5.95 (1H, m), 5.34-5.06 (2H, m), 3.37-2.92 (5H, m), 2.70 (1H, m), 2.38-2.15 (3H, m), 1.94-1.52 (2H, m)

(e) [R]-2-(6-Methoxyquinolin-4-yl)oxirane

A solution of 6-methoxyquinoline-4-carboxylic acid (10g) in dichloromethane was heated under reflux with oxalyl chloride (5ml) and dimethylformamide (2 drops) for
 10 1 hour and evaporated to dryness. ~~The residue, in dichloromethane (100ml) was treated~~
 with a 2M solution of trimethylsilyldiazomethane in hexane (50ml) and stirred at room temperature for 18 hours. 5M Hydrochloric acid (150ml) was added and the solution was stirred at room temperature for 3 hours. It was basified with sodium carbonate solution, extracted with ethyl acetate and chromatographed on silica gel eluting with ethyl acetate-
 15 hexane to give the chloromethyl ketone (4.2g). A batch of the chloromethyl ketone (20g) was reduced with (+)-B-chlorodiisopinocampheylborane (40g) in dichloromethane (400ml) at room temperature for 18 hours followed by treatment with diethanolamine (30 g) for 3 hours. The product was chromatographed on silica gel eluting with ethyl acetate-hexane to give the chloroalcohol (16.8g), which was dissolved in tetrahydrofuran (100
 20 ml) and reacted with sodium hydroxide (2.6g) in water (13ml) for 1.5 hours. The reaction mixture was evaporated to dryness and chromatographed on silica gel eluting with ethyl acetate - hexane to give the title compound as a solid (10.4 g) (84% ee by chiral HPLC). Recrystallisation from ether-pentane gave mother-liquor (7.0 g) (90% ee).


MS (+ve ion electrospray) m/z 202 (MH⁺)

25 The absolute stereochemistry was defined to be (R) by an NMR study on the Mosher's esters derived from the product obtained by reaction with 1-t-butylpiperazine.

(f) Title compounds

To an enantiomeric mixture of 3-fluoro-4-[[[1,2,3]thiadiazolo[5,4-b]pyridine-6-ylmethyl)-amino]-piperidine (0.33 mmole) in DMF (2 mL) is added LiClO₄ (0.33
 30 mmole), K₂CO₃ (0.66 mmole), and 6-methoxy-4-(R)-oxiranylquinoline (0.33 mmole). The reaction is heated at 90 °C for 18 h, cooled to RT and concentrated under reduced pressure. The residue is partitioned between ethyl acetate and H₂O, and the layers are separated. The aqueous phase is further extracted with ethyl acetate, and the combined
 35 organic extracts are dried over MgSO₄ and concentrated under reduced pressure. The residue is purified by flash chromatography on silica gel to give the title compounds.

Example 2. Preparation of 5-[(3*S*,4*R*)-3-fluoro-1-[(*R*)-2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]piperidin-4-ylamino}methyl]-benzo[1,2,3]thiadiazole and 5-[(3*R*,4*S*)-3-fluoro-1-[(*R*)-2-hydroxy-2-(6-methoxyquinolin-4-yl)-ethyl]piperidin-4-ylamino}methyl]-benzo[1,2,3]thiadiazole



(a) Benzo[1,2,3]thiadiazole-5-carboxaldehyde

10 Methyl benzo[1,2,3]thiadiazole-5-carboxylate (0.291g) in THF (2ml) was treated dropwise with lithium aluminium hydride (1M in THF, 0.71ml) at 0 °C.

After 1.5h, additional lithium aluminium hydride (0.1ml) was added. After 0.5h the mixture was treated with 8% sodium hydroxide, ethyl acetate and sodium sulphate, filtered and evaporated. The resulting crude alcohol was dissolved in dichloromethane
15 (3ml) and stirred with manganese (II) oxide overnight. Filtration and evaporation of solvent gave the aldehyde .

(b) Title compounds

These are prepared analogously to Example 1 using aldehyde (a).


20

Biological Activity

The MIC (µg/ml) of test compounds against various organisms may be determined: *S. aureus* Oxford, *S. aureus* WCUH29, *S. pneumoniae* 1629, *S. pneumoniae* N1387, *S. pneumoniae* ERY 2, *E. faecalis* 1, *H. influenzae* Q1, *M. catarrhalis* 1502.

Claims

1. a compound of formula (I) or a pharmaceutically acceptable derivative thereof:



wherein:

one of Z^1 , Z^2 , Z^3 , Z^4 and Z^5 is N, one is CR^{1a} and the remainder are CH, or one or two of Z^1 , Z^2 , Z^3 , Z^4 and Z^5 are independently CR^{1a} and the remainder are CH;

R^1 and R^{1a} are independently hydrogen; hydroxy; (C₁₋₆)alkoxy optionally substituted by (C₁₋₆)alkoxy, amino, piperidyl, guanidino or amidino any of which is optionally N-substituted by one or two (C₁₋₆)alkyl, acyl or (C₁₋₆)alkylsulphonyl groups, CONH₂, hydroxy, (C₁₋₆)alkylthio, heterocyclylthio, heterocyclyloxy, arylthio, aryloxy, acylthio, acyloxy or (C₁₋₆)alkylsulphonyloxy; (C₁₋₆)alkoxy-substituted (C₁₋₆)alkyl; halogen; (C₁₋₆)alkyl; (C₁₋₆)alkylthio; trifluoromethyl; trifluoromethoxy; nitro; azido; acyl; acyloxy; acylthio; (C₁₋₆)alkylsulphonyl; (C₁₋₆)alkylsulphoxide; arylsulphonyl; arylsulphoxide or an amino, piperidyl, guanidino or amidino group optionally N-substituted by one or two (C₁₋₆)alkyl, acyl or (C₁₋₆)alkylsulphonyl groups;

provided that when Z^1 , Z^2 , Z^3 , Z^4 and Z^5 are CR^{1a} or CH, then R^1 is not hydrogen;


R^2 is hydrogen, or (C₁₋₄)alkyl or (C₂₋₄)alkenyl optionally substituted with 1 to 3 groups selected from:

amino optionally substituted by one or two (C₁₋₄)alkyl groups; carboxy; (C₁₋₄)alkoxycarbonyl; (C₁₋₄)alkylcarbonyl; (C₂₋₄)alkenyloxycarbonyl; (C₂₋₄)alkenylcarbonyl; aminocarbonyl wherein the amino group is optionally substituted by hydroxy, (C₁₋₄)alkyl, hydroxy(C₁₋₄)alkyl, aminocarbonyl(C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₁₋₄)alkylsulphonyl, trifluoromethylsulphonyl, (C₂₋₄)alkenylsulphonyl, (C₁₋₄)alkoxycarbonyl, (C₁₋₄)alkylcarbonyl, (C₂₋₄)alkenyloxycarbonyl or (C₂₋₄)alkenylcarbonyl; cyano; tetrazolyl; 2-oxo-oxazolidinyl optionally substituted by R^{10} ; 3-hydroxy-3-cyclobutene-1,2-dione-4-yl; 2,4-thiazolidinedione-5-yl; tetrazol-5-ylaminocarbonyl; 1,2,4-triazol-5-yl optionally substituted by R^{10} ; 5-oxo-1,2,4-oxadiazol-

3-yl; halogen; (C₁₋₄)alkylthio; trifluoromethyl; hydroxy optionally substituted by (C₁₋₄)alkyl, (C₂₋₄)alkenyl, (C₁₋₄)alkoxycarbonyl, (C₁₋₄)alkylcarbonyl, (C₂₋₄)alkenyloxycarbonyl, (C₂₋₄)alkenylcarbonyl; oxo; (C₁₋₄)alkylsulphonyl; (C₂₋₄)alkenylsulphonyl; or (C₁₋₄)aminosulphonyl wherein the amino group is optionally substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl;

R³ is in the 2-, 3- or 4-position and is trifluoromethyl or is in the 2-position and is oxo; or R³ is in the 3-position and is fluorine or amino wherein the amino group is optionally substituted by: hydroxy; (C₁₋₆)alkylsulphonyl; trifluoromethylsulphonyl; (C₂₋₆)alkenylsulphonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenylcarbonyl; (C₁₋₆)alkoxycarbonyl; (C₂₋₆)alkenyloxycarbonyl; (C₁₋₆)alkyl; or (C₂₋₆)alkenyl; wherein a (C₁₋₆)alkyl or (C₂₋₆)alkenyl moiety may be optionally substituted with up to 2 groups R¹² independently selected from:
halogen; (C₁₋₆)alkylthio; trifluoromethyl; cyano; carboxy; tetrazolyl; 2-oxo-oxazolidinyl; 3-hydroxy-3-cyclobutene-1,2-dione-4-yl; 2,4-thiazolidinedione-5-yl; tetrazol-5-ylaminocarbonyl; 1,2,4-triazol-5-yl optionally substituted by R¹⁰; or 5-oxo-1,2,4-oxadiazol-3-yl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; (C₂₋₆)alkenylcarbonyl; hydroxy optionally substituted by (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenylcarbonyl or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl, (C₂₋₆)alkenyl; amino optionally mono- or disubstituted by (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl, (C₂₋₆)alkenylcarbonyl, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylsulphonyl, (C₂₋₆)alkenylsulphonyl or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl;
in addition when R³ is disubstituted with a hydroxy or amino containing substituent and carboxy containing substituent these may together form a cyclic ester or amide linkage, respectively;

R⁴ is a group -U-R⁵ where
U is selected from CO, SO₂ and CH₂ and
R⁵ is an optionally substituted bicyclic carbocyclic or heterocyclic ring system (A):



containing up to four heteroatoms in each ring in which

ring (a) is aromatic or non aromatic;
X¹ is C when part of an aromatic ring or CR¹⁴ when part of a non aromatic ring;

X^2 is N, NR^{13} , O, $S(O)_x$, CO or CR^{14} when part of an aromatic or non-aromatic ring or may in addition be $CR^{14}R^{15}$ when part of a non aromatic ring;

X^4 is N, NR^{13} , O, $S(O)_x$, CO or CR^{14} ;

X^3 and X^5 are independently N or C;

- 5 Y^1 is a 1 to 3 atom linker group each atom of which is independently selected from N, NR^{13} , O, $S(O)_x$, CO and CR^{14} when part of an aromatic or non-aromatic ring or may additionally be $CR^{14}R^{15}$ when part of a non aromatic ring,

Y^2 is a 2 or 3 atom linker group completing an aromatic ring, each atom of Y^2 being independently selected from N, NR^{13} , O, $S(O)_x$, CO and CR^{14} ;

- 10 each of R^{14} and R^{15} is independently selected from: H; (C₁₋₄)alkylthio; halo; carboxy(C₁₋₄)alkyl; halo(C₁₋₄)alkoxy; halo(C₁₋₄)alkyl; (C₁₋₄)alkyl; (C₂₋₄)alkenyl; (C₁₋₄)alkoxycarbonyl; formyl; (C₁₋₄)alkylcarbonyl; (C₂₋₄)alkenyloxycarbonyl; (C₂₋₄)alkenylcarbonyl; (C₁₋₄)alkylcarbonyloxy; (C₁₋₄)alkoxycarbonyl(C₁₋₄)alkyl; hydroxy; hydroxy(C₁₋₄)alkyl; mercapto(C₁₋₄)alkyl; (C₁₋₄)alkoxy; nitro; cyano; carboxy; amino or
15 aminocarbonyl optionally substituted as for corresponding substituents in R^3 ; (C₁₋₄)alkylsulphonyl; (C₂₋₄)alkenylsulphonyl; or aminosulphonyl wherein the amino group is optionally mono- or di-substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl; aryl; aryl(C₁₋₄)alkyl; aryl(C₁₋₄)alkoxy or

R^{14} and R^{15} may together represent oxo;

- 20 each R^{13} is independently H; trifluoromethyl; (C₁₋₄)alkyl optionally substituted by hydroxy, (C₁₋₆)alkoxy, (C₁₋₆)alkylthio, carboxy, halo or trifluoromethyl; (C₂₋₄)alkenyl; aryl; aryl (C₁₋₄)alkyl; arylcarbonyl; heteroarylcarbonyl; (C₁₋₄)alkoxycarbonyl; (C₁₋₄)alkylcarbonyl; formyl; (C₁₋₆)alkylsulphonyl; or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₄)alkoxycarbonyl, (C₁₋₄)alkylcarbonyl, (C₂₋₄)alkenyloxycarbonyl, (C₂₋₄)alkenylcarbonyl, (C₁₋₄)alkyl or (C₂₋₄)alkenyl and
25 optionally further substituted by (C₁₋₄)alkyl or (C₂₋₄)alkenyl;

n is 0 or 1;

- 30 each x is independently 0, 1 or 2

A is NR^{11} , O or CR^6R^7 and B is NR^{11} , O, SO_2 or CR^8R^9 and wherein:

- each of R^6 , R^7 , R^8 and R^9 is independently selected from: hydrogen; (C₁₋₆)alkoxy; (C₁₋₆)alkylthio; halo; trifluoromethyl; azido; (C₁₋₆)alkyl; (C₂₋₆)alkenyl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; (C₂₋₆)alkenylcarbonyl; hydroxy, amino or aminocarbonyl optionally substituted as for
35 corresponding substituents in R^3 ; (C₁₋₆)alkylsulphonyl; (C₂₋₆)alkenylsulphonyl; or

aminosulphonyl wherein the amino group is optionally substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl;

or when n=1 R⁶ and R⁸ together represent a bond and R⁷ and R⁹ are as above defined;
or R⁶ and R⁷ or R⁸ and R⁹ together represent oxo;

5

provided that:

when A is NR¹¹, B is not NR¹¹ or O;

when A is CO, B is not CO, O or SO₂;

when n is 0 and A is NR¹¹, CR⁸R⁹ can only be CO;

10

when A is CR⁶R⁷ and B is SO₂, n is 0;

when n is 0, B is not NR¹¹ or O or R⁸ and R⁹ are not optionally substituted hydroxy or amino;

when A is O, B is not NR¹¹, O, SO₂ or CO and n=1; and

when A-B is CR⁷=CR⁹, n is 1

15

R¹⁰ is selected from (C₁₋₄)alkyl; (C₂₋₄)alkenyl and aryl any of which may be optionally substituted by a group R¹² as defined above; carboxy; aminocarbonyl wherein the amino group is optionally substituted by hydroxy, (C₁₋₆)alkyl, (C₂₋₆)alkenyl, (C₁₋₆)alkylsulphonyl, trifluoromethylsulphonyl, (C₂₋₆)alkenylsulphonyl, (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl or (C₂₋₆)alkenylcarbonyl and optionally further substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl; (C₁₋₆)alkylsulphonyl; trifluoromethylsulphonyl; (C₂₋₆)alkenylsulphonyl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; (C₂₋₆)alkenyloxycarbonyl; and (C₂₋₆)alkenylcarbonyl; and

25


R¹¹ is hydrogen; trifluoromethyl, (C₁₋₆)alkyl; (C₂₋₆)alkenyl; (C₁₋₆)alkoxycarbonyl; (C₁₋₆)alkylcarbonyl; or aminocarbonyl wherein the amino group is optionally substituted by (C₁₋₆)alkoxycarbonyl, (C₁₋₆)alkylcarbonyl, (C₂₋₆)alkenyloxycarbonyl, (C₂₋₆)alkenylcarbonyl, (C₁₋₆)alkyl or (C₂₋₆)alkenyl and optionally further substituted by (C₁₋₆)alkyl or (C₂₋₆)alkenyl;

30


or where one of R³ and R⁶, R⁷, R⁸ or R⁹ contains a carboxy group and the other contains a hydroxy or amino group they may together form a cyclic ester or amide linkage.

35 2. A compound according to claim 1 wherein Z⁵ is CH or N, Z³ is CH or CF and Z¹, Z² and Z⁴ are each CH, or Z¹ is N, Z³ is CH or CF and Z², Z⁴ and Z⁵ are each CH.

3. A compound according to any preceding claim wherein R^1 is methoxy, amino(C_{3-5})alkyloxy or guanidino(C_{3-5})alkyloxy.
4. A compound according to any preceding claim wherein R^2 is hydrogen,
5 carboxymethyl, hydroxyethyl, aminocarbonylmethyl, ethoxycarbonylmethyl, ethoxycarbonylallyl or carboxyallyl.
5. A compound according to any preceding claim wherein R^3 is CF_3 , fluoro, oxo or amino unsubstituted or substituted by (C_{1-6})alkyl or (C_{2-6})alkenyl.
10
6. A compound according to any preceding claim wherein n is 0, A-B is $CHOH-CH_2$, $NR^{11}-CH_2$ or $NR^{11}-CO$ and R^{11} is hydrogen or (C_{1-4})alkyl.
7. A compound according to any preceding claim wherein U is CH_2 and R^5 is an
15 aromatic heterocyclic ring (A) having 1-4 heteroatoms of which one is N or NR^{13} , R^{13} is H if in ring (a) or in addition (C_{1-4})alkyl, R^{14} and R^{15} are independently selected from hydrogen, halo, hydroxy, (C_{1-4})alkyl, (C_{1-4})alkoxy, trifluoromethoxy, nitro, cyano, aryl(C_{1-4})alkoxy and (C_{1-4})alkylsulphonyl.
8. A compound according to any of claims 1 to 6 wherein R^5 is 4,6-difluoro-indol-2-yl, 1H-pyrrolo[2,3-b]-pyridin-2-yl, 1H-pyrrolo[3,2-b]-pyridin-2-yl, 8-hydroxy-quinolin-2-yl, quinoxalin-2-yl, benzimidazol-2-yl, benzo[1,2,3]-thiadiazol-5-yl, benzothiophen-2-yl, 4,6-difluoro-1H-benzimidazol-2-yl, benzothiazole-5-yl or 3-(R)-2,3-dihydro-[1,4]dioxino[2,3-b]pyridin-3-yl.
20
25
9. A method of treatment of bacterial infections in mammals, particularly in man, which method comprises the administration to a mammal in need of such treatment an effective amount of a compound according to claim 1.
10. The use of a compound according to claim 1, in the manufacture of a medicament
30 for use in the treatment of bacterial infections in mammals.
11. A pharmaceutical composition comprising a compound according to claim 1, and a pharmaceutically acceptable carrier.
12. A process for preparing compounds according to claim 1, which process
35 comprises:
reacting a compound of formula (IV) with a compound of formula (V):



(IV)




(V)

wherein n is as defined in formula (I); $Z^{1'}$, $Z^{2'}$, $Z^{3'}$, $Z^{4'}$, $Z^{5'}$, $R^{1'}$ and $R^{3'}$ are Z^1 , Z^2 , Z^3 , Z^4 , Z^5 , R^1 and R^3 as defined in formula (I) or groups convertible thereto;
 Q^1 is $NR^{2'}R^{4'}$ or a group convertible thereto wherein $R^{2'}$ and $R^{4'}$ are R^2 and R^4 as defined in formula (I) or groups convertible thereto and Q^2 is H or $R^{3'}$ or Q^1 and Q^2 together form an optionally protected oxo group;
 and X and Y may be the following combinations:

- (i) X is $A'-COW$, Y is H and n is 0;
- (ii) X is $CR^6=CR^8R^9$, Y is H and n is 0;
- (iii) X is oxirane, Y is H and n is 0;
- (iv) X is $N=C=O$ and Y is H and n is 0;
- (v) one of X and Y is CO_2R^Y and the other is $CH_2CO_2R^X$;
- (vi) X is CHR^6R^7 and Y is $C(=O)R^9$;
- (vii) X is $CR^7=PR^{Z_3}$ and Y is $C(=O)R^9$ and $n=1$;
- (viii) X is $C(=O)R^7$ and Y is $CR^9=PR^{Z_3}$ and $n=1$;
- (ix) Y is COW and X is $NHR^{11'}$, NCO or $NR^{11'}COW$ and $n=0$ or 1 or when $n=1$ X is COW and Y is $NHR^{11'}$, NCO or $NR^{11'}COW$;
- (x) X is $C(O)=R^6$ and Y is $NHR^{11'}$ or X is $NHR^{11'}$ and Y is $C(=O)R^8$ and $n=1$;
- (xi) X is $NHR^{11'}$ and Y is CR^8R^9W and $n=1$;
- (xii) X is CR^6R^7W and Y is $NHR^{11'}$ or OH and $n=1$;
- (xiii) X is $CR^6R^7SO_2W$ and Y is H and $n=0$;
- (xiv) X is W or OH and Y is CH_2OH and $n=1$;
- (xv) X is $NHR^{11'}$ and Y is SO_2W or X is $NR^{11'}SO_2W$ and Y is H, and $n=0$;
- (xvi) X is $NR^{11'}COCH_2W$ or $NR^{11'}SO_2CH_2W$ and Y is H and $n=0$;
- (xvii) X is W and Y is $CONHR^{11'}$;

in which W is a leaving group, e.g. halo or imidazolyl; R^X and R^Y are (C_{1-6}) alkyl; R^Z is aryl or (C_{1-6}) alkyl; A' and $NR^{11'}$ are A and NR^{11} as defined in formula (I), or groups convertible thereto; and oxirane is:



- wherein R^6 , R^8 and R^9 are as defined in formula (I);
and thereafter optionally or as necessary converting Q^1 and Q^2 to NR^2R^4 ; converting A' ,
 Z^1 , Z^2 , Z^3 , Z^4 , Z^5 , R^1 , R^2 , R^3 , R^4 and NR^{11} to A , Z^1 , Z^2 , Z^3 , Z^4 , Z^5 , R^1 , R^2 , R^3 ,
 R^4 and NR^{11} ; converting A-B to other A-B, interconverting R^1 , R^2 , R^3 and/or R^4 ,
5 and/or forming a pharmaceutically acceptable derivative thereof.

ABSTRACT

Compounds

Aminopiperidine derivatives and pharmaceutically acceptable derivatives thereof useful in methods of treatment of bacterial infections in mammals, particularly in man.

5

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.